

Transition to longitudinal instability of detonation waves is generically associated with Hopf bifurcation to time-periodic galloping solutions

BENJAMIN TEXIER* AND KEVIN ZUMBRUN†

December 22, 2008

Abstract

We show that transition to longitudinal instability of strong detonation solutions of reactive compressible Navier–Stokes equations is generically associated with Hopf bifurcation to nearby time-periodic “galloping”, or “pulsating”, solutions, in agreement with physical and numerical observation. The analysis is by pointwise semigroup techniques introduced by the authors and collaborators in previous works.

Contents

1	Introduction	2
1.1	The reacting Navier-Stokes equations	3
1.2	Assumptions	4
1.3	Coordinatizations	4
1.4	Strong detonations	5
1.5	Structure of the equations and the profiles	8
1.6	The Evans function	8
1.7	Results	10
1.7.1	Stability	10
1.7.2	Transition from stability to instability	10
1.7.3	Nonlinear instability	12
1.8	Discussion and open problems	12
2	Strong detonations	13

*Université Paris Diderot (Paris 7), Institut de Mathématiques de Jussieu, UMR CNRS 7586; texier@math.jussieu.fr: Research of B.T. was partially supported under NSF grant number DMS-0505780.

†Indiana University, Bloomington, IN 47405; kzumbrun@indiana.edu: Research of K.Z. was partially supported under NSF grants no. DMS-0300487 and DMS-0801745.

3	Resolvent kernel and Green function bounds	17
3.1	Laplace transform	17
3.1.1	The limiting, constant-coefficient equations	19
3.1.2	Low-frequency behaviour of the normal modes	21
3.1.3	Description of the essential spectrum	23
3.1.4	Gap Lemma and dual basis	25
3.1.5	Duality relation and forward basis	28
3.1.6	The resolvent kernel	31
3.1.7	The Evans function	33
3.2	Inverse Laplace transform	33
3.2.1	Pointwise Green function bounds	33
3.2.2	Convolution bounds	37
4	Stability: Proof of Theorem 1.12	37
4.1	Linearized stability criterion	37
4.2	Auxiliary energy estimate	38
4.3	Nonlinear stability	38
5	Bifurcation: Proof of Theorem 1.14	42
5.1	The perturbation equations	42
5.2	Coordinatization	43
5.3	Poincaré return map	43
5.4	Lyapunov-Schmidt reduction	44
5.4.1	Pointwise cancellation estimate	45
5.4.2	Reduction	49
5.4.3	Bifurcation	50
6	Nonlinear instability: Proof of Theorem 1.15	51

1 Introduction

Motivated by physical and numerical observations of time-oscillatory “galloping” or “pulsating” instabilities of detonation waves [MT, BMR, FW, MT, AIT, AT, F1, F2, KS], we study Hopf bifurcation of viscous detonation waves, or traveling-wave solutions of the reactive compressible Navier–Stokes equations. This extends a larger program begun in [Z1, LyZ1, LyZ2, JLW, LRTZ] toward the dynamical study of viscous combustion waves using Evans function/inverse Laplace transform techniques introduced in the context of viscous shock waves [GZ, ZH, ZS, Z1, MaZ3], continuing the line of investigation initiated in [TZ1, TZ2, SS, TZ3] on bifurcation/transition to instability.

It has long been observed that transition to instability of detonation waves occurs in certain predictable ways, with the archetypal behavior in the case of longitudinal,

or one-dimensional instability being transition from a steady planar progressing wave $U(x, t) = \bar{U}(x_1 - st)$ to a galloping, or time-periodic planar progressing wave $\bar{U}(x_1 - st, t)$, where $\bar{U}$ is periodic in the second coordinate, and in the case of transverse, or multi-dimensional instability, transition to more complicated “spinning” or “cellular behavior”; see [KS, TZ1, TZ2], and references therein.

The purpose of this paper is, restricting to the one-dimensional case, to establish this principle rigorously, arguing from first principles from the physical equations that *transition to longitudinal instability of detonation waves is generically associated with Hopf bifurcation to time-periodic galloping solutions*.

1.1 The reacting Navier-Stokes equations

The single-species reactive compressible Navier–Stokes equations, in Lagrangian coordinates, appear as [Ch]

$$(1.1) \quad \begin{cases} \partial_t \tau - \partial_x u = 0, \\ \partial_t u + \partial_x p = \partial_x (\nu \tau^{-1} \partial_x u), \\ \partial_t E + \partial_x (pu) = \partial_x (qd\tau^{-2} \partial_x z + \kappa \tau^{-1} \partial_x T + \nu \tau^{-1} u \partial_x u), \\ \partial_t z + k\phi(T)z = \partial_x (d\tau^{-2} \partial_x z), \end{cases}$$

where $\tau > 0$ denotes specific volume, u velocity, $E > 0$ total specific energy, and $0 \leq z \leq 1$ mass fraction of the reactant.

The variable

$$U := (\tau, u, E, z) \in \mathbb{R}^4$$

depend on time $t \in \mathbb{R}_+$, position $x \in \mathbb{R}$, and parameters ν, κ, d, k, q , where $\nu > 0$ is a viscosity coefficient, $\kappa > 0$ and $d > 0$ are respectively coefficients of heat conduction and species diffusion, $k > 0$ represents the rate of the reaction, and q is the heat release parameter, with $q > 0$ corresponding to an exothermic reaction and $q < 0$ to an endothermic reaction.

In (1.1), $T = T(\tau, e, z) > 0$ represents temperature, $p = p(\tau, e, z)$ pressure, where the internal energy $e > 0$ is defined through the relation

$$E = e + \frac{1}{2}u^2 + qz.$$

In (1.1), we assume a simple one-step, one-reactant, one-product reaction



where ϕ is an ignition function. More realistic reaction models are described in [GS2].

In the variable U , after the shift

$$x \rightarrow x - st, \quad s \in \mathbb{R},$$

the system (1.1) takes the form of a system of differential equations

$$(1.2) \quad \partial_t U + \partial_x(F(U)) = \partial_x(B(U)\partial_x U) + G(U),$$

where

$$F := \begin{pmatrix} -u \\ p \\ pu \\ 0 \end{pmatrix} - s(\varepsilon)U, \quad G := \begin{pmatrix} 0 \\ 0 \\ 0 \\ -k\phi(T)z \end{pmatrix},$$

and

$$B := \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & \nu\tau^{-1} & 0 & 0 \\ \kappa\tau^{-1}\partial_\tau T & -\kappa u\tau^{-1}\partial_e T + \nu\tau^{-1}u & \kappa\tau^{-1}\partial_e T & \kappa\tau^{-1}(\partial_z T - q\partial_e T) + qd\tau^{-2} \\ 0 & 0 & 0 & d\tau^{-2} \end{pmatrix}.$$

The characteristic speeds of the first-order part of (1.1), i.e., the eigenvalues of $\partial_U F(U)$, are

$$(1.3) \quad \left\{ \underbrace{-s - \sigma, -s, -s + \sigma}_{\text{fluid eigenvalues}}, \underbrace{-s}_{\text{reactive eigenvalue}} \right\},$$

where σ , the sound speed of the gas, is

$$\sigma := (p\partial_e p - \partial_\tau p)^{\frac{1}{2}} = \tau^{-1}(\Gamma(\Gamma + 1)e)^{\frac{1}{2}}.$$

1.2 Assumptions

We make the following assumptions:

Assumption 1.1. *We assume a reaction-independent ideal gas equation of state,*

$$p = \Gamma\tau^{-1}e, \quad T = c^{-1}e,$$

where $c > 0$ is the specific heat constant and Γ is the Gruneisen constant.

Assumption 1.2. *The ignition function ϕ is smooth; it vanishes identically for $T \leq T_i$, and is strictly positive for $T > T_i$.*

1.3 Coordinatizations

We let

$$w := (u, E, z) \in \mathbb{R}^3, \quad v := (\tau, u, E) \in \mathbb{R}^3.$$

Then we have the coordinatizations

$$U = (v, z) = (\tau, w).$$

In particular, Assumption 1.1 implies that in the (τ, w) coordinatization, B takes the block-diagonal form

$$B = \begin{pmatrix} 0 & 0 \\ 0 & b \end{pmatrix},$$

where b is full rank for all values of the parameters and U ; the system (1.2) in (τ, w) coordinates is

$$\begin{cases} \partial_t \tau - s \partial_x \tau - J \partial_x w = 0, \\ \partial_t w + \partial_x f(\tau, w) = \partial_x (b(\tau, w) \partial_x w) + g(w), \end{cases}$$

with the notation

$$(1.4) \quad J := \begin{pmatrix} 1 & 0 & 0 \end{pmatrix}, \quad f := \begin{pmatrix} p \\ pu \\ 0 \end{pmatrix} - sw, \quad g := \begin{pmatrix} 0 \\ 0 \\ -k\phi(T)z \end{pmatrix}.$$

In the (v, z) coordinatization, the system (1.2) takes the form

$$\begin{cases} \partial_t v + \partial_x f^\sharp(v, z) = \partial_x (b_1^\sharp(v) \partial_x v + b_2^\sharp(v) \partial_x z) \\ \partial_t z - s \partial_x z + k\phi(T)z = \partial_x (d\tau^{-2} \partial_x z). \end{cases}$$

where the flux is $f^\sharp = (-u - s\tau, p - su, pu - sE)$, and, under Assumption 1.1, the diffusion matrices are

$$b_1^\sharp = \begin{pmatrix} 0 & 0 & 0 \\ 0 & \nu\tau^{-1} & 0 \\ 0 & \tau^{-1}(\nu - \kappa c^{-1})u & \kappa\tau^{-1}c^{-1} \end{pmatrix}, \quad b_2^\sharp = \begin{pmatrix} 0 \\ 0 \\ q(d\tau^{-2} - \kappa\tau^{-1}) \end{pmatrix}.$$

Note that, in the (v, z) coordinatization, the first component is a conservative variable, in the sense that $\partial_t v$ is a perfect derivative, hence

$$(1.5) \quad \int_{\mathbb{R}} (v(x, t) - v(x, 0)) dx \equiv 0,$$

for $v(t) - v(0) \in W^{2,1}(\mathbb{R})$.

1.4 Strong detonations

We prove in this article stability and bifurcation results for viscous strong detonations of (1.1), defined as follows:

Definition 1.3. *A one-parameter, right-going family of viscous strong detonations is a family $\{\bar{U}^\varepsilon\}_{\varepsilon \in \mathbb{R}}$ of smooth stationary solutions of (1.2), associated with speeds $s(\varepsilon)$, $s(\varepsilon) > 0$, model parameters $(\nu, \kappa, d, k, q)(\varepsilon)$ and ignition function ϕ^ε , with $\bar{U}^\varepsilon, \phi^\varepsilon, (s, \nu, \kappa, d, k, q)(\varepsilon)$ depending smoothly on ε in $L^\infty \times L^\infty \times \mathbb{R}^6$, satisfying*

$$(1.6) \quad \bar{U}^\varepsilon(x, t) = \bar{U}^\varepsilon(x), \quad \lim_{x \rightarrow \pm\infty} \bar{U}^\varepsilon(x) = U_\pm^\varepsilon,$$




Figure 1: Characteristic speeds for strong detonations.

connecting a burned state on the left to an unburned state on the right,

$$(1.7) \quad z_-^\varepsilon \equiv 0, \quad z_+^\varepsilon \equiv 1,$$

with a temperature on the burned side above ignition temperature

$$(1.8) \quad T_-^\varepsilon > T_i,$$

and satisfying the Lax characteristic conditions

$$(1.9) \quad \sigma_- := \sigma(U_-^\varepsilon) > s > \sigma_+ := \sigma(U_+^\varepsilon),$$

uniformly in ε .

Consider a standing wave (1.6), $U = (\tau, u, E, z)$, solution of (1.2), with endstates $U_\pm = (\tau_\pm, u_\pm, E_\pm, z_\pm)$. It satisfies the linear constraint

$$-s(\tau - \tau_-) = u - u_-,$$

the system of ordinary differential equations

$$(1.10) \quad \begin{cases} \nu\tau^{-1}u' = p - su - (p - su)_-, \\ \kappa\tau^{-1}c^{-1}E' + \tau^{-1}(\nu - \kappa c^{-1})uu' = pu - sE - (pu - sE)_- \\ \quad + (\kappa\tau^{-1}c^{-1} - d\tau^{-2})qy, \\ z' = y, \\ d\tau^{-2}y' = -sy + k\phi(T)z, \end{cases}$$

and the Rankine-Hugoniot relations

$$(1.11) \quad \begin{cases} -s(\tau_+ - \tau_-) = u_+ - u_-, \\ (p - su)_+ = (p - su)_-, \\ (pu - sE)_+ = (pu - sE)_-, \\ y_{\pm} = 0, \\ \phi(T_{\pm})z_{\pm} = 0, \end{cases}$$

expressing the fact that $(u_{\pm}, E_{\pm}, 0, z_{\pm})$ are rest points of (1.10).

From (1.9) and (1.11), we note that the right endstate of a strong detonation satisfies

$$(1.12) \quad \phi(T_+^{\varepsilon}) = 0,$$

which, by Assumption 1.2, implies also

$$(1.13) \quad \phi'(T_+^{\varepsilon}) = 0.$$

Lemma 1.4. *Under Assumptions 1.1, 1.2, if $q > 0$ and s is large enough with respect to q , then for any $z_+ \in (0, 1]$, there exists an open subset $\mathcal{O}_-$ in $\mathbb{R}^3$, such that any left endstate $U_- = (v_-, 0)$ with $v_- \in \mathcal{O}_-$ satisfies (1.8) and (1.9), and is associated with a right endstate $U_+ = (v_+, z_+)$ satisfying $T_+ < T_i$, (1.9) and (1.11).*

The existence of strong detonations was proved by Gasser and Szmolyan [GS1] for small dissipation coefficients ν, κ and d . We restrict throughout the article to strong detonations with left endstates as in the above Lemma.

Remark 1.5. *In the small-heat-release limit $q \rightarrow 0$, the equations in (y, z) (in system (1.10)) are decoupled from the fluid equations; in particular, strong detonations converge to ordinary nonreacting gas-dynamical shocks of standard Lax type, the existence of which was first proved by Gilbarg [G].*

A consequence of Lemma 1.4 is that strong detonations converge exponentially to their endstates, a key fact of the subsequent stability and bifurcation analysis.

Corollary 1.6. *Under Assumptions 1.1, 1.2, let $\{\bar{U}^{\varepsilon}\}_{\varepsilon}$ be a family of viscous strong detonations. There exist $C, \eta_0 > 0$, such that, for $k \geq 0$ and $j \in \{0, 1\}$,*

$$(1.14) \quad \begin{aligned} |\partial_{\varepsilon}^j \partial_x^k (\bar{U}^{\varepsilon} - U_-^{\varepsilon})(x)| &\leq C e^{-\eta_0 |x|}, & x < 0, \\ |\partial_{\varepsilon}^j \partial_x^k (\bar{U}^{\varepsilon} - U_+^{\varepsilon})(x)| &\leq C e^{-\eta_0 |x|}, & x > 0. \end{aligned}$$

In particular, $|(\bar{U}^{\varepsilon})'(x)| \leq C e^{-\eta_0 |x|}$, for all x .

Remark 1.7. *In the ZND limit, strong detonations are transverse orbits of (1.10), a result proved in Section 3.6 of [LyZ2], after [GS1].*

Lemma 1.4 and Corollary 1.6 are proved in Section 2.

1.5 Structure of the equations and the profiles

System (1.2), seen as a system in τ, w , satisfies

- (A1) the convection terms in the equation in τ are linear in (τ, w) ;
- (A2) the diffusion matrix b is positive definite.

For strong detonation waves, the convection terms in (1.1) satisfy

- (H1) The convection coefficient $s(\varepsilon)$ in the evolution equation in τ is nonzero, uniformly in ε .
- (H2) The spectrum of $\partial_U F$, given in (1.3), is real, simple, and nonzero, uniformly in ε .

System (1.2) satisfies the Kawashima dissipativity condition

- (H3) For all ε , for all $\xi \in \mathbb{R}$,

$$\Re \sigma(i\xi \partial_U F(U_\pm^\varepsilon) - \xi^2 B(U_\pm^\varepsilon) + \partial_U G(U_\pm^\varepsilon)) \leq -\frac{\theta \xi^2}{1 + \xi^2},$$

at the endstates $U_\pm^\varepsilon$ of a family of strong detonations. In (H3), σ denotes spectrum of a matrix, and $\theta > 0$ is independent of ξ and ε . To verify (H3), it suffices, by a classical result of [ShK], to check that (1.2) has a symmetrizable hyperbolic-parabolic structure, and that the genuine coupling condition holds. These conditions are coordinates-independent, and easily checked in (τ, u, e) coordinates.

Finally, the assumption

- (H4) Considered as connecting orbits of (1.10), $\bar{U}^\varepsilon$ lie in a smooth one-dimensional manifold of solutions of (1.10), obtained as a transversal intersection of the unstable manifold at U_-^ε and the stable manifold at U_+^ε ,

holds in the ZND limit, as stated in Remark 1.7. Under (H4), in a vicinity of $\bar{U}^\varepsilon$, the set of stationary solutions of (1.2) with limits $U_\pm^\varepsilon$ at $\pm\infty$ is a smooth one-dimensional manifold, given by $\{\bar{U}^\varepsilon(\cdot - c), \quad c \in \mathbb{R}\}$, and the associated speed $\varepsilon \rightarrow s(\varepsilon)$ is smooth.

Conditions (A1)-(A2), (H0)-(H4) are the assumptions of [TZ3] (where $G \equiv 0$), themselves a strengthened version of the assumptions of [MaZ3].

1.6 The Evans function

A central object in the study of stability of traveling waves is the *Evans function* $D(\varepsilon, \cdot)$ (precisely defined in Section 3.1.7), a Wronskian of solutions of the eigenvalue

equation $(L(\varepsilon) - \lambda)U = 0$ decaying at plus or minus spatial infinity [AGJ]¹, where the linearized operator L is defined as

$$(1.15) \quad L(\varepsilon) := -\partial_x(A \cdot) + \partial_x(B(\bar{U}^\varepsilon)\partial_x \cdot) + \partial_U G(\bar{U}^\varepsilon),$$

with the notation

$$(1.16) \quad A := -\partial_U F(\bar{U}^\varepsilon) + (\partial_U B(\bar{U}^\varepsilon) \cdot)(\bar{U}^\varepsilon)'.$$

Recall the important result of [LyZ2]:

Proposition 1.8 ([LyZ2], Theorem 4). *Under Assumptions 1.1 and 1.2, let $\{\bar{U}^\varepsilon\}_\varepsilon$ be a one-parameter family of viscous strong detonation waves satisfying (H4).*

For all ε , the associated Evans function has a zero of multiplicity one at $\lambda = 0$:

$$D(\varepsilon, 0) = 0, \quad \text{and} \quad D'(\varepsilon, 0) \neq 0.$$

Proof. By translational invariance, $D(\varepsilon, 0) = 0$, for all ε . Generalizing similar results known for shock waves [GZ, ZS], there was established in [Z1, LyZ1, LyZ2] the fundamental relation

$$(1.17) \quad D'(\varepsilon, 0) = \gamma \delta.$$

In (1.17), γ is a coefficient given as a Wronskian of solutions of the linearized traveling-wave ODE about $\bar{U}$; transversality corresponds to $\gamma \neq 0$. In (1.17), δ is the Lopatinski determinant

$$\delta := \det \begin{pmatrix} r_1^- & r_2^- & r_4^- & (\tau_+ - \tau_- & u_+ - u_- & E_+ - E_- + q)^{\text{tr}} \end{pmatrix},$$

(where r_j^- denote the eigenvectors of $\partial_U F(U_-^\varepsilon)$ associated with outgoing eigenvalues, F as in (1.2), and tr denotes transverse matrix²) determining hyperbolic stability of the Chapman–Jouget (square wave) approximation modeling the detonation as a shock discontinuity. Hyperbolic stability corresponds to $\delta \neq 0$. See [Z1, LyZ1, LyZ2, JLW] for further discussion. By (H4), $\gamma \neq 0$, while $\delta \neq 0$ by direct calculation comparing to the nonreactive (shock-wave) case. $\square$

Remark 1.9. *The vectors r_1^- , r_2^- and r_4^- correspond to outgoing modes to the left of $x = 0$, see Section 3.1.2 and Figure 4. (The fluid modes r_j^- , $1 \leq j \leq 3$, are ordered as usual by increasing characteristic speeds: $-s - \sigma_- < -s < 0 < -s + \sigma_-$, so that r_3^- is incoming.)*

¹For applications of the Evans function to stability of viscous shock and detonation waves, see, e.g., [AGJ, GZ, ZS, Z1, LyZ1, LyZ2, LRTZ].

²This notation will be used throughout the article.

1.7 Results

Let X and Y be two Banach spaces, and consider a traveling wave $\bar{U}$ solution of a general evolution equation.

Definition 1.10. A traveling wave $\bar{U}$ is said to be $X \rightarrow Y$ linearly orbitally stable if, for any solution $\tilde{U}$ of the linearized equations about $\bar{U}$ with initial data in X , there exists a phase shift δ , such that $|\tilde{U}(\cdot, t) - \delta(t)\bar{U}'(\cdot)|_Y$ is bounded for $0 \leq t \leq \infty$.

It is said to be $X \rightarrow Y$ linearly asymptotically orbitally stable if it is $X \rightarrow Y$ linearly orbitally stable and if moreover $|\tilde{U}(\cdot, t) - \delta(t)\bar{U}'(\cdot)|_Y \rightarrow 0$ as $t \rightarrow \infty$.

Definition 1.11. A traveling wave $\bar{U}$ is said to be $X \rightarrow Y$ nonlinearly orbitally stable if, for each $\delta > 0$, for any solution $\tilde{U}$ of the nonlinear equations with $|\tilde{U}(\cdot, 0) - \bar{U}|_X$ sufficiently small, there exists a phase shift δ , such that $|\tilde{U}(\cdot, t) - \bar{U}(\cdot - \delta(t), t)|_Y \leq \delta$ for $0 \leq t \leq \infty$.

It is said to be $X \rightarrow Y$ nonlinearly asymptotically orbitally stable if it is $X \rightarrow Y$ nonlinearly orbitally stable and if moreover $|\tilde{U}(\cdot, t) - \bar{U}(\cdot - \delta(t), t)|_Y \rightarrow 0$ as $t \rightarrow \infty$.

1.7.1 Stability

Our first result, generalizing that of [LRTZ] in the artificial viscosity case, is a characterization of linearized stability and a sufficient condition for nonlinear stability, in terms of an Evans function condition.

Theorem 1.12. Under Assumptions 1.1, 1.2, let $\{\bar{U}^\varepsilon\}_\varepsilon$ be a one-parameter family of viscous strong detonation waves.

For all ε , $\bar{U}^\varepsilon$ is $L^1 \cap L^p \rightarrow L^p$ linearly orbitally stable if and only if, for all ε ,

$$(1.18) \quad \text{the only zero of } D(\varepsilon, \cdot) \text{ in } \Re \lambda \geq 0 \text{ is a simple zero at the origin.}$$

If (1.18) holds, $\bar{U}^\varepsilon$ is $L^1 \cap H^3 \rightarrow L^1 \cap H^3$ linearly and nonlinearly orbitally stable, and $L^1 \cap H^3 \rightarrow L^p \cap H^3$ asymptotically orbitally stable, for $p > 1$, with

$$(1.19) \quad |\tilde{U}^\varepsilon(\cdot, t) - \bar{U}^\varepsilon(\cdot - \delta(t))|_{L^p} \leq C|\tilde{U}_0^\varepsilon - \bar{U}^\varepsilon|_{L^1 \cap H^3}(1+t)^{-\frac{1}{2}(1-\frac{1}{p})},$$

where $\tilde{U}^\varepsilon$ is the solution of (1.2) issued from $\tilde{U}_0^\varepsilon$, for some $\delta(\cdot)$ satisfying

$$\begin{aligned} |\delta(t)| &\leq C|\tilde{U}_0^\varepsilon - \bar{U}^\varepsilon|_{L^1 \cap H^3}, \\ |\dot{\delta}(t)| &\leq C|\tilde{U}_0^\varepsilon - \bar{U}^\varepsilon|_{L^1 \cap H^3}(1+t)^{-\frac{1}{2}}. \end{aligned}$$

1.7.2 Transition from stability to instability

Theorem 1.13. Under Assumptions 1.1, 1.2, let $\{\bar{U}^\varepsilon\}_\varepsilon$ be a one-parameter family of viscous strong detonation waves satisfying (H4).

Assume that the family of equations (1.2) and profiles $\bar{U}^\varepsilon$ undergoes transition to instability at $\varepsilon = 0$ in the sense that $\bar{U}^\varepsilon$ is linearly stable for $\varepsilon < 0$ and linearly unstable for $\varepsilon > 0$.

Then, one or more pair of nonzero complex conjugate eigenvalues of $L(\varepsilon)$ move from the stable (negative real part) to the neutral or unstable (nonnegative real part) half-plane as ε passes from negative to positive through $\varepsilon = 0$, while $\lambda = 0$ remains a simple root of $D(\varepsilon, \cdot)$ for all ε .

That is, transition to instability is associated with a Hopf-type bifurcation in the spectral configuration of the linearized operator about the wave.

Proof of Theorem 1.13. By Theorem 1.12, transition from stability to instability must occur through the passage of a root of the Evans function from the stable half-plane to the neutral or unstable half-plane. However, Proposition 1.8 implies that D has a zero of multiplicity one at the origin, for all ε , and so no root can pass through the origin. It follows that transition to instability, if it occurs, must occur through the passage of one or more nonzero complex conjugate pairs $\lambda = \gamma \pm i\tau$, $\tau \neq 0$, from the stable half-plane ($\gamma < 0$ for $\varepsilon < 0$) to the neutral or unstable half-plane ($\gamma \geq 0$ for $\varepsilon \geq 0$). $\square$

Our third result and the main object of this paper is to establish, under appropriate nondegeneracy conditions, that the spectral Hopf bifurcation configuration described in Theorem 1.13 is realized at the nonlinear level as a genuine bifurcation to time-periodic solutions.

Given $k \in \mathbb{N}$ and a weight function $\omega > 0$, define the Sobolev space and associated norm

$$(1.20) \quad H_\omega^k := \{f \in \mathcal{S}'(\mathbb{R}), \omega^{\frac{1}{2}}f \in H^k(\mathbb{R})\}, \quad \|f\|_{H_\omega^k} := \|\omega^{\frac{1}{2}}f\|_{H^k}.$$

Let $\omega \in C^2$ be a growing weight function such that, for some $\theta_0 > 0$, $C > 0$, for all x, y ,

$$(1.21) \quad \begin{cases} 1 \leq \omega(x) \leq e^{\theta_0(1+|x|^2)^{\frac{1}{2}}}, \\ |\omega'(x)| + |\omega''(x)| \leq C\omega(x), \\ \omega(x) \leq C\omega(x-y)\omega(y). \end{cases}$$

Theorem 1.14. *Under Assumptions 1.1, 1.2, let $\{\bar{U}^\varepsilon\}_\varepsilon$ be a family of viscous strong detonation waves satisfying (H4).*

Assume that the family of equations (1.2) and profiles $\bar{U}^\varepsilon$ undergoes transition from linear stability to linear instability at $\varepsilon = 0$.

Moreover, assume that this transition is associated with passage of a single complex conjugate pair of eigenvalues of $L(\varepsilon)$, $\lambda_\pm(\varepsilon) = \gamma(\varepsilon) \pm i\tau(\varepsilon)$ through the imaginary axis, satisfying

$$(1.22) \quad \gamma(0) = 0, \quad \tau(0) \neq 0, \quad d\gamma/d\varepsilon(0) \neq 0.$$

Then, given a growing weight ω satisfying (1.21) with θ_0 sufficiently small, for $r \geq 0$ sufficiently small and $C > 0$ sufficiently large, there are C^1 functions $r \rightarrow \varepsilon(r)$, $r \rightarrow T(r)$, with $\varepsilon(0) = 0$, $T(0) = 2\pi/\tau(0)$, and a C^1 family of time-periodic solutions $\tilde{U}^r(x, t)$ of (1.2) with $\varepsilon = \varepsilon(r)$, of period $T(r)$, with

$$(1.23) \quad C^{-1}r \leq \|\tilde{U}^r - \bar{U}^\varepsilon\|_{H_\omega^2} \leq Cr,$$

Up to translation in x, t , these are the only time-periodic solutions nearby in $\|\cdot\|_{H_\omega^2}$ with period $T \in [T_0, T_1]$ for any fixed $0 < T_0 < T_1 < +\infty$.

That is, transition to linear instability of viscous strong detonation waves is “generically” (in the sense of (1.22)) associated with Hopf bifurcation to time-periodic galloping solutions, as asserted in the title of this paper.

The choices $\omega \equiv 1$ and $\omega = e^{\theta_0(1+|x|^2)^{\frac{1}{2}}}$ are allowed in (1.21), as well as $\omega = (1 + |x|^2)^p$, for any real $p > 0$. In Theorem 1.14, we need, in particular, $\theta_0 < \eta_0$, where η_0 is as in Corollary 1.6, so that the spatial localization given by (1.23) is less precise than the spatial localization of the background profile $\bar{U}^\varepsilon$. The smallness condition on θ_0 is described in Remark 5.9.

1.7.3 Nonlinear instability

We complete our discussion with the following straightforward result verifying that the exchange of linear stability described in Theorem 1.14, as expected, corresponds to an exchange of nonlinear stability as well, the new assertion being nonlinear *instability* for $\varepsilon > 0$.

Theorem 1.15. *Under the assumptions of Theorem 1.14, the viscous strong detonation waves $\bar{U}^\varepsilon$ undergo a transition at $\varepsilon = 0$ from nonlinear orbital stability to instability; that is, $\bar{U}^\varepsilon$ is nonlinearly orbitally stable for $\varepsilon < 0$ and unstable for $\varepsilon > 0$.*

1.8 Discussion and open problems

This analysis in large part concludes the one-dimensional program set out in [TZ2]. However, a very interesting remaining open problem is to determine linearized and nonlinear stability of the bifurcating time-periodic solutions, in the spirit of Section 4.3. For a treatment in the shock wave case with semilinear viscosity, see [BeSZ]. Likewise, it would be very interesting to carry out a numerical investigation of the spectrum of the linearized operator about detonation waves with varying physical parameters, as done in [LS, KS] in the inviscid ZND setting, but using the viscous methods of [Br1, Br2, BrZ, BDG, HuZ] to treat the full reacting Navier–Stokes equations, in order to determine the physical bifurcation boundaries.

Other interesting open problems are the extension to multi-dimensional (spinning or cellular) bifurcations, as carried out for artificial viscosity systems in [TZ2], and to the case of weak detonations (analogous to the case of undercompressive viscous shocks; see [HZ, RZ, LRTZ]).

The strong detonation structure considerably simplifies both stability and bifurcation arguments over what was done in [LRTZ]. We remark that, at the expense of further complication, nonlinear stability of general (time-independent) combustion waves, including also weak detonations and strong or weak deflagrations, may be treated by a combination of the pointwise arguments of [LRTZ] and [RZ].

We remark finally that the restriction to a scalar reaction variable is for simplicity only. Indeed, the results of this article (as well as the results of the article by Lyng and Zumbrun [LyZ2] from which it draws) are independent from the dimension of the reactive equation, so long as the reaction satisfies an assumption of exponential decay of space-independent states (with temperature at $-\infty$ above the ignition temperature).

Plan of the paper. Lemma 1.4 and Corollary 1.6 are proved in Section 2. We give a detailed description of the low-frequency behavior of the resolvent kernel for the linearized equations in Section 3, following [MaZ3]. In Section 4, we prove Theorem 1.12, while Section 5 is devoted to the proof of Theorem 1.14. Finally, in Section 6, we prove Theorem 1.15.

Acknowledgement. Thanks to Björn Sandstede and Arnd Scheel for their interest in this work and for stimulating discussions on spatial dynamics and bifurcation in the absence of a spectral gap. Thanks to Gregory Lyng for pointing out reference [Ch]. B.T. thanks Indiana University for their hospitality during the collaborative visit in which the analysis was carried out. B.T. and K.Z. separately thank the Ecole Polytechnique Fédérale de Lausanne for their hospitality during two visits in which a substantial part of the analysis was carried out.

2 Strong detonations

Proof of Lemma 1.4. Let U_- be a given left endstate, with $z_- = 0$, satisfying (1.8) and (1.9). We look for a right endstate U_+ , with $z_+ \in (0, 1]$, that satisfies (1.11), (1.9), and $T_+ < T_i$. We note that (1.11)(i) determines u_+ and that $T_+ < T_i$ entails (1.11)(v).

The Rankine-Hugoniot relations in the (τ_+, p_+) plane are

$$\begin{cases} p = -s^2\tau + c_1 & \text{(R),} \\ p = (c_0 - s\tau(1 + \Gamma^{-1}))^{-1}(c_2 + sqz_+ + \frac{1}{2}s^3\tau^2 - s^2c_0\tau) & \text{(H),} \end{cases}$$

where (R) is the Rayleigh line, corresponding to (1.11)(ii), (H) the Hugoniot curve, corresponding to (1.11)(iii), and where

$$c_0 := u_- + s\tau_-, \quad c_1 := p_- + s^2\tau_-, \quad c_2 := (p_-u_- - sE_-) + \frac{1}{2}c_0^2s$$

depend on parameters U_- and s . The temperature and Lax constraints for both enstates are

$$\begin{cases} \tau_+ p_+ < c\Gamma T_i < \tau_- p_- & (\text{T})_{\pm}, \\ \tau_+^{-1} p_+ < (\Gamma + 1)^{-1} s^2 < \tau_-^{-1} p_- & (\text{L})_{\pm}. \end{cases}$$

We restrict to left endstates satisfying in the large s regime

$$(2.1) \quad \tau_- = O(1), \quad p_- = 2s^2\Gamma^{-1}\tau_- + \tilde{p}_-, \quad u_- = s\tilde{u}_-,$$

with $\tilde{u}_- = O(1)$ and $\tilde{p}_- = O(1)$. Under (2.1), conditions $(\text{T})_-$ and $(\text{L})_-$ are satisfied as soon as s is large enough. The Hugoniot curve takes the form

$$p_H = (\tilde{u}_- + \tau_- - (1 + \Gamma^{-1})\tau)^{-1} \left(\frac{1}{2}s^3(\tau - (1 + 2\Gamma^{-1})\tau_-)(\tau - (1 - 2\Gamma^{-1})\tau_- - 2\tilde{u}_-) + sqz_+ \right).$$

Assume that $\tilde{u}_-$ is such that

$$(2.2) \quad \frac{\tau_-}{1 + \Gamma^{-1}} < (1 - 2\Gamma^{-1})\tau_- + 2\tilde{u}_- < (1 + 2\Gamma^{-1})\tau_-.$$

For any such $\tilde{u}_-$, any given τ_- and any $q > 0$, if s is large enough then, for any $z_+ \in (0, 1]$, the Hugoniot curve has two zeros $\underline{\tau} < \bar{\tau}$, with asymptotic expansions

$$(2.3) \quad \underline{\tau} = (1 - 2\Gamma^{-1})\tau_- + 2\tilde{u}_- + O(s^{-2}).$$

$$(2.4) \quad \bar{\tau} = (1 + 2\Gamma^{-1})\tau_- - s^{-2} \frac{\tilde{p}_-\tilde{u}_- + qz_+}{2\Gamma^{-1}\tau_- - \tilde{u}_-} + O(s^{-3}).$$

If s is large, by (2.2), $\tau_0 < \underline{\tau} < \bar{\tau}$, where $\tau_0 := c_0 s^{-1}(1 + \Gamma^{-1})^{-1}$ is the pole of (H).

The Rayleigh line and the Hugoniot curve have at least one intersection point to the right of τ_0 if

$$p_R(\bar{\tau}) < 0 < p_R(\underline{\tau}).$$

Under (2.2), the inequality $0 < p_R(\underline{\tau})$ holds, and $p_R(\bar{\tau}) < 0$ holds as well if in addition

$$(2.5) \quad \tilde{p}_- < -\frac{\tilde{p}_-\tilde{u}_- + qz_+}{2\Gamma^{-1}\tau_- - \tilde{u}_-}.$$

Let τ_+ be an intersection point of (R) and (H) to the right of τ_0 . Condition $(\text{T})_+$ is satisfied if

$$(2.6) \quad \tau_+ = (1 + 2\Gamma^{-1})\tau_- + s^{-2}\tilde{\tau}_+ + O(s^{-3}),$$

with

$$(2.7) \quad (1 + 2\Gamma^{-1})\tau_-(\tilde{p}_- - \tilde{\tau}_+) < c\Gamma T_i.$$

Condition (L)₊ is satisfied if

$$(2.8) \quad (1 + 2\Gamma^{-1})\tau_- < (1 + (1 + \Gamma)^{-1})\tau_+,$$

which holds under (2.6), if s is large. We plug the ansatz (2.6) in the equation $p_H = p_R$, to find

$$(2.9) \quad \tilde{\tau}_+ = \frac{\Gamma\tilde{p}_-}{(1 + 2\Gamma^{-1})\tau_-} ((1 + \Gamma^{-1})(1 + 2\Gamma^{-1}) - 1) + \frac{\Gamma qz_+}{(1 + 2\Gamma^{-1})\tau_-}.$$

The intersection point τ_+ is an admissible right specific volume if $p_H(\tau_+) > 0$ and $p_R(\tau_+) > 0$. These inequalities holds if

$$(2.10) \quad \underline{\tau} < \tau_+ < (\alpha + 1)\tau_- + s^{-2}\tilde{p}_-.$$

The inequalities (2.5), (2.7) and (2.10) are constraints on τ_- , $\tilde{p}_-$, and $\tilde{u}_-$. The lower bound on τ_+ in (2.10) is satisfied in the regime (2.1) if s is large. If we let

$$\tilde{p}_- = \frac{-2\Gamma qz_+}{\tau_-} + O(s^{-1}),$$

then (2.5) holds. Finally, if τ_- satisfies

$$1 < \frac{1}{4\tau_-} (3 + 2\Gamma^{-1} - (1 + 2\Gamma^{-1})\tau_-) < 1 + \frac{cT_i}{qz_+}.$$

then the upper bound in (2.10) and (2.7) hold as well.

The Rayleigh line (R), the Hugoniot curve (R) and the temperature (T) and Lax (L) constraints are pictured on Figure 2. The black dots represent the intersection points of (R) and (H). Note that (L) and (R) imply $\tau_- < \tau_+$ for a strong detonation, so that only the intersection point to the right to τ_- is admissible. (The other intersection point corresponds to a deflagration, see for instance [LyZ2], Section 1.4.) $\square$

Proof of Corollary 1.6. Rewrite (1.10) as $U' = \mathfrak{F}(\varepsilon, U)$. Let $U_\pm^\varepsilon$ be the endstates of a family of strong detonations. The linearized equations at $U_\pm^\varepsilon$ are governed by matrices

$$(2.11) \quad \partial_U \mathfrak{F}(\varepsilon, \bar{U}_\pm^\varepsilon) = \begin{pmatrix} a_\pm^f & * \\ 0 & a_\pm^r \end{pmatrix}.$$

The block triangular structure is a consequence of Assumption 1.1, (1.7), (1.12), and (1.13). Under Assumption 1.1, the eigenvalues λ of $a_\pm^f$

$$a_\pm^f := \begin{pmatrix} \nu\tau^{-1} & 0 \\ \tau^{-1}(\nu - \kappa c^{-1})u & \kappa c^{-1}\tau^{-1} \end{pmatrix} \begin{pmatrix} \partial_u p - s - s^{-1}\partial_\tau p & \partial_e p \\ u(\partial_u p - s^{-1}\partial_\tau p) + p & u\partial_e p - s \end{pmatrix},$$




Figure 2: The Rankine-Hugoniot, Lax and temperature conditions.

satisfy

$$(2.12) \quad \lambda^2 + (s\kappa c^{-1}\tau_{\pm}^{-1} + s^{-1}\nu\tau_{\pm}^{-1}(s^2 + (\partial_{\tau}p)_{\pm}))\lambda + \kappa c^{-1}\nu\tau_{\pm}^{-2}(s^2 - \sigma_{\pm}^2) = 0.$$

The Lax condition (1.9) implies that the center subspace on both sides is trivial, that the eigenvalues of a_{-}^f have opposite signs, and that the eigenvalues of a_{+}^f are negative. The eigenvalues λ of

$$a_{\pm}^r := \begin{pmatrix} -sd^{-1} & kd^{-1}\phi(T_{\pm}) \\ 1 & 0 \end{pmatrix}$$

satisfy

$$d\lambda^2 + s\lambda - k\phi(T_{\pm}) = 0.$$

They are non zero and have distinct signs on the $-\infty$ side. On the $+\infty$ side, there is one negative eigenvalue, and a one-dimensional kernel. In particular, U_{-}^{ε} is a hyperbolic rest point of the linearized traveling-wave ordinary differential equation, which implies (1.14)(i) with $j = 0$, by standard ODE estimates. However, the linearized traveling-wave equations at U_{+} have a one-dimensional center subspace, which a priori precludes exponential decay (1.14).

From Lemma 1.4, if $U_{-} \in \mathcal{O}$, then the system (1.10) has a line of equilibria that goes through U_{+}^{ε} . Any center manifold of (1.10) at U_{+}^{ε} contains all equilibria, so by dimension count it must consist of equilibria. Therefore, the 4-dimensional stable

center manifold at U_+^ε consists (again by dimension count) of the union of the stable manifolds of all equilibria. Since solutions off of stable center manifold do not stay for all time in small vicinity of center manifold, any traveling-wave orbit must lie on the center-stable manifold, so lies on the stable manifold of some equilibrium. Exponential decay, (1.14)(ii), $j = 0$, now follows by the stable manifold theorem.

To prove (1.14) with $j = 1$, consider now the traveling-wave ODE in $(U, \partial_\varepsilon U)$. The rest points satisfy

$$(2.13) \quad \mathfrak{F}(\varepsilon, U) = 0, \quad \partial_\varepsilon \mathfrak{F}(\varepsilon, U) + \partial_U \mathfrak{F}(\varepsilon, U) \partial_\varepsilon U = 0.$$

The kernel of $\partial_U \mathfrak{F}(\varepsilon, U_+^\varepsilon)$ being one-dimensional, (2.13) has a two-dimensional manifold of solutions. Let $(U_+^\varepsilon, V_+^\varepsilon)$ be such a rest point. The linearized equations at $(U_+^\varepsilon, V_+^\varepsilon)$ are governed by matrices

$$\begin{pmatrix} \partial_U \mathfrak{F}(\varepsilon, U_+^\varepsilon) & 0 \\ * & \partial_U \mathfrak{F}(\varepsilon, U_+^\varepsilon) \end{pmatrix},$$

where the bottom left entry depends on second derivatives of $\mathfrak{F}$. In particular, the linearized equations have a two-dimensional center subspace. We can thus argue as above that any center manifold consists entirely of equilibria, and that (1.14)(ii) holds with $j = 1$. The proof of (1.14)(i) with $j = 1$ is similar. $\square$

3 Resolvent kernel and Green function bounds

The linearized equations about a traveling wave $\bar{U}^\varepsilon$ solution of (1.2) are

$$(3.1) \quad \partial_t U = L(\varepsilon)U,$$

where $L(\varepsilon)$ is defined in (1.15). The coefficients of $L(\varepsilon)$ are asymptotically constant at $\pm\infty$. Let $L_\pm(\varepsilon)$ be the associated constant-coefficient, limiting operators:

$$L_\pm(\varepsilon) := -A_\pm \partial_x + B_\pm \partial_x^2 + G_\pm,$$

with the notation $A_\pm := \partial_U F(U_\pm^\varepsilon)$, $B_\pm := B(U_\pm^\varepsilon)$, $G_\pm := \partial_U G(U_\pm^\varepsilon)$. Let $L(\varepsilon)^*$ denote the dual operator of $L(\varepsilon)$. Its associated constant-coefficient, limiting operators are $L_\pm(\varepsilon)^* = A_\pm^* \partial_x + B_\pm^* \partial_x^2 + G_\pm^*$.

3.1 Laplace transform

Consider the Laplace transform of the linearized equations,

$$(3.2) \quad (L(\varepsilon) - \lambda)U = 0, \quad \lambda \in \mathbb{C}, \quad x \in \mathbb{R}, \quad U(\varepsilon, x, \lambda) \in \mathbb{C}^4.$$

Equation (3.2) can be cast as a first-order ordinary differential system in $\mathbb{R}^7$,

$$(3.3) \quad W' = \mathbb{A}(\varepsilon, \lambda)W, \quad \lambda \in \mathbb{C}, \quad x \in \mathbb{R}, \quad W(\varepsilon, x, \lambda) \in \mathbb{C}^7,$$

where the limits $\mathbb{A}_\pm$ of $\mathbb{A}$ at $\pm\infty$ are given by

$$(3.4) \quad \mathbb{A}_\pm := \begin{pmatrix} s^{-1}\lambda & 0 & -s^{-1}Jb_\pm^{-1} \\ 0 & 0 & b_\pm^{-1} \\ s^{-1}\lambda\partial_\tau f|_\pm & \lambda - \partial_w g|_\pm & (\partial_w f|_\pm - s^{-1}\partial_\tau f|_\pm J)b_\pm^{-1} \end{pmatrix},$$

where $|_\pm$ denotes evaluation at $U_\pm^\varepsilon$, $b_\pm := b(U_\pm^\varepsilon)$, and J is defined in (1.4).

Considered as an operator in $L^2(\mathbb{R}; \mathbb{C}^4)$, L is closed, with domain H^2 dense in L^2 . Similarly, for all λ , the operator

$$\frac{d}{dx} - \mathbb{A}(\lambda) : H^1(\mathbb{R}; \mathbb{C}^7) \subset L^2(\mathbb{R}; \mathbb{C}^7) \rightarrow L^2(\mathbb{R}; \mathbb{C}^7)$$

is closed and densely defined.

The following straightforward Lemma gives a correspondence between (3.2) and (3.3).

Lemma 3.1. *Let $\lambda \in \mathbb{C}$ and $f = (f_1, f_2) \in L^2(\mathbb{R}; \mathbb{C}^1 \times \mathbb{C}^3)$. If the equation*

$$(3.5) \quad (L - \lambda)U = f$$

has a solution $U =: (\tau, w) \in H^2(\mathbb{R}; \mathbb{C}^1 \times \mathbb{C}^3)$, then $W := (\tau, w, bw') \in H^1(\mathbb{R}; \mathbb{C}^7)$ satisfies

$$(3.6) \quad W' = \mathbb{A}(\lambda)W + F,$$

with $F = (f_1, 0, f_2) \in L^2(\mathbb{R}; \mathbb{C}^7)$. Conversely, let $F = (f_1, 0, f_2) \in L^2(\mathbb{R}; \mathbb{C}^7)$ and $\lambda \in \mathbb{C}$. If $W = (w_1, w_2) \in H^1(\mathbb{R}; \mathbb{C}^7)$ satisfies (3.6), then a solution in $H^2(\mathbb{R}; \mathbb{C}^4)$ to (3.5) with $f = (f_1, f_2)$ is given by $U = w_1$.

Similarly, the dual eigenvalue equation

$$(3.7) \quad (L(\varepsilon)^* - \lambda)\tilde{U} = 0, \quad \lambda \in \mathbb{C}, \quad y \in \mathbb{R}, \quad \tilde{U}(\varepsilon, y, \lambda) \in \mathbb{C}^4,$$

can be cast as

$$(3.8) \quad \tilde{W}' = \tilde{\mathbb{A}}(\varepsilon, \lambda)\tilde{W}, \quad \lambda \in \mathbb{C}, \quad y \in \mathbb{R}, \quad \tilde{W}(\varepsilon, y, \lambda) \in \mathbb{C}^7,$$

where the limits $\tilde{\mathbb{A}}_\pm$ of $\tilde{\mathbb{A}}$ at $\pm\infty$ are given by

$$(3.9) \quad \tilde{\mathbb{A}}_\pm := \begin{pmatrix} -s^{-1}\lambda & 0 & s^{-1}\partial_\tau f|_\pm^{\text{tr}} b_\pm^{\text{tr}-1} \\ 0 & 0 & b_\pm^{\text{tr}-1} \\ s^{-1}\lambda J^{\text{tr}} & \lambda - \partial_w g|_\pm^{\text{tr}} & -(\partial_w f|_\pm^{\text{tr}} + s^{-1}J^{\text{tr}}\partial_\tau f|_\pm^{\text{tr}})b_\pm^{\text{tr}-1} \end{pmatrix}.$$

A correspondence between (3.7) and (3.8) holds, as in Lemma 3.1.

3.1.1 The limiting, constant-coefficient equations

Associated with (3.2) and (3.7) are the limiting, constant-coefficient eigenvalue equations

$$(3.10) \quad (L_{\pm}(\varepsilon) - \lambda)U = 0,$$

and

$$(3.11) \quad (L_{\pm}(\varepsilon)^* - \lambda)\tilde{U} = 0.$$

Definition 3.2 (Normal modes). *We call normal modes the solutions (λ, U) of equations (3.10) and dual normal modes the solutions $(\lambda, \tilde{U})$ of equations (3.11).*

Associated with (3.3) and (3.8) are the limiting, constant-coefficient differential equations

$$(3.12) \quad W' = \mathbb{A}_{\pm}(\varepsilon, \lambda)W,$$

and

$$(3.13) \quad \tilde{W}' = \tilde{\mathbb{A}}_{\pm}(\varepsilon, \lambda)\tilde{W},$$

where $\mathbb{A}_{\pm}$ and $\tilde{\mathbb{A}}_{\pm}$ are defined in (3.4) and (3.9).

There is a correspondence between solutions of (3.10) and solutions of (3.12):

Lemma 3.3. *If (λ_0, U) , $U =: (\tau, w)$, is a normal mode, then $W := (\tau, w, bw')$ solves (3.12) at $\lambda = \lambda_0$. Conversely, if $W = (w_1, w_2) \in \mathbb{C}^4 \times \mathbb{C}^3$ solves (3.12) at $\lambda = \lambda_0$, then (λ_0, w_1) is a normal mode. In particular,*

(i) *Eigenvalues μ of $\mathbb{A}_{\pm}$ satisfy*

$$(3.14) \quad \det(-\mu A_{\pm} + \mu^2 B_{\pm} + G_{\pm} - \lambda) = 0,$$

and associated eigenvectors, satisfying $\mathbb{A}_{\pm}(\lambda)W = \mu W$, have the form $W = (U, w_2) \in \mathbb{C}^4 \times \mathbb{C}^3$, with

$$(3.15) \quad U \in \ker(-\mu A_{\pm} + \mu^2 B_{\pm} + G_{\pm} - \lambda), \quad U =: (\tau, w), \quad w_2 := \mu b_{\pm} w,$$

(ii) *Normal modes (λ, U) satisfy*

$$(3.16) \quad U = \sum_j e^{x\mu_j^{\pm}(\lambda)} U_j^{\pm}(x, \lambda),$$

where the $\mu_j^{\pm}$ are eigenvalues of $\mathbb{A}_{\pm}$, and the $U_j^{\pm}$ are polynomials in x .

The correspondence between (3.11) and (3.13) is similar. In particular, eigenvalues $\tilde{\mu}$ of $\tilde{\mathbb{A}}_{\pm}$ satisfy

$$(3.17) \quad \det(\tilde{\mu}A_{\pm}^* + \tilde{\mu}^2B_{\pm}^* + G_{\pm}^* - \lambda) = 0,$$

associated eigenvectors, satisfying $\tilde{\mathbb{A}}_{\pm}(\lambda)\tilde{W} = \tilde{\mu}\tilde{W}$, have the form $\tilde{W} = (\tilde{U}, \tilde{w}_2) \in \mathbb{C}^4 \times \mathbb{C}^3$, with

$$(3.18) \quad \tilde{U} \in \ker(\tilde{\mu}A_{\pm}^* + \tilde{\mu}^2B_{\pm}^* + G_{\pm}^* - \lambda), \quad \tilde{U} =: (\tilde{\tau}, \tilde{w}), \quad \tilde{w}_2 := \mu b_{\pm}^{\text{tr}} \tilde{w},$$

and dual normal modes satisfy

$$(3.19) \quad \tilde{U} = \sum_j e^{y\tilde{\mu}_j^{\pm}(\lambda)} \tilde{U}_j^{\pm}(y, \lambda),$$

where the $\tilde{\mu}_j^{\pm}$ are eigenvalues of $\tilde{\mathbb{A}}_{\pm}$, and the $\tilde{U}_j^{\pm}$ are polynomials in y .

If $\tilde{\mu}(\lambda)$ is an eigenvalue of $\tilde{\mathbb{A}}_{\pm}(\lambda)$, then $\overline{\tilde{\mu}(\lambda)} = -\mu(\bar{\lambda})$, where $\mu(\bar{\lambda})$ is some eigenvalue of $\mathbb{A}_{\pm}(\bar{\lambda})$. The matrices $A_{\pm}, B_{\pm}$ and $G_{\pm}$ having real coefficients, the complex conjugate of $\mu(\bar{\lambda})$ is an eigenvalue of $\mathbb{A}_{\pm}(\lambda)$. We can thus relate the solutions of (3.14) and (3.17) by

$$\tilde{\mu}(\lambda) = -\mu(\lambda).$$

Note that $z_+^{\varepsilon} = 1$, $\phi(T_+^{\varepsilon}) = 0$, and $\phi'(T_+^{\varepsilon}) = 0$ imply that the v derivative of the coupling reaction term $k\phi(T)z$ vanishes when evaluated at $U_{\pm}^{\varepsilon}$. In particular, in (v, z) coordinates,

$$A_{\pm} = \begin{pmatrix} \partial_v f_{|\pm}^{\sharp} & \partial_z f_{|\pm}^{\sharp} \\ 0 & -s \end{pmatrix}, B_{\pm} = \begin{pmatrix} b_{1|\pm}^{\sharp} & b_{2|\pm}^{\sharp} \\ 0 & d \end{pmatrix}, G_{\pm} = \begin{pmatrix} 0 & 0 \\ 0 & -k\phi_{\pm} \end{pmatrix},$$

with the notation of Section 1.3, $|\pm$ denoting evaluation at $U_{\pm}^{\varepsilon}$, and $\phi_{\pm} := \phi(T_{\pm}^{\varepsilon})$, so that $\phi_+ = 0$, while by (1.8) and Assumption 1.2, $\phi_- > 0$. This triangular structure of the matrix $-\mu A_{\pm} + \mu^2 B_{\pm} + G_{\pm}$ allows a simple description of the solutions of (3.14). Indeed, (3.14), a polynomial, degree four equation in λ , splits into the linear equation

$$(3.20) \quad \mu s + \mu^2 d - k\phi_{\pm} - \lambda = 0,$$

and the degree three equation

$$(3.21) \quad \det(-\mu \partial_v f_{|\pm}^{\sharp} + \mu^2 b_{1|\pm}^{\sharp} - \lambda) = 0.$$

By inspection, (3.20) is quadratic in μ , while (3.21) is degree five in μ . Thus, the four solutions $\lambda(\mu)$ of (3.14) correspond to seven eigenvalues $\mu(\lambda)$ of $\mathbb{A}(\lambda)$.

3.1.2 Low-frequency behaviour of the normal modes

We describe here the behaviour of the normal modes in a small ball $B(0, r) := \{\lambda \in \mathbb{C}, |\lambda| < r\}$.

Definition 3.4 (Slow modes, fast modes). *We call slow mode at $\pm\infty$ any family of normal modes*

$$\{(\lambda, U(\lambda))\}_{\lambda \in B(0, r)}, \quad \text{for some } r > 0,$$

such that, in (3.16), $\mu_j^\pm(0) = 0$, for all j . Normal modes which are not slow are called fast modes. We define similarly slow dual modes and fast dual modes, using (3.19).

The solutions of (3.20) are

$$(3.22) \quad \mu_4^\pm = \frac{1}{2d}(-s + (s^2 + 4d(\lambda + k\phi_\pm))^{\frac{1}{2}}),$$

$$(3.23) \quad \mu_5^\pm = -\frac{1}{2d}(s + (s^2 + 4d(\lambda + k\phi_\pm))^{\frac{1}{2}});$$

they depend analytically on λ (in the case of μ_+^4 and μ_5^+ , this is ensured by $s > 0$, assumed in Definition 1.3), and satisfy, for λ in a neighborhood of the origin,

$$(3.24) \quad \mu_4^+ = s^{-1}\lambda - s^{-3}d\lambda^2 + O(\lambda^3), \quad \mu_4^- > 0, \quad \mu_5^\pm < 0.$$

Note that the inequality $\mu_4^- > 0$ is a consequence of $\phi_- > 0$. By (3.18), the eigenvector of $\tilde{\mathbb{A}}_+$ that is associated with $-\mu_4^+$ is

$$(3.25) \quad L_4^+ = \begin{pmatrix} \underline{\ell}_4^+ \\ \mu_4^+ b_+^{\text{tr}} \underline{\ell}_4^+ \end{pmatrix} \in \mathbb{C}^4 \times \mathbb{C}^3, \quad \underline{\ell}_4^+(0) = \ell_4^+,$$

where

$$(3.26) \quad \ell_4^+ := \begin{pmatrix} 0 & 0 & 0 & 1 \end{pmatrix}^{\text{tr}}$$

is the reactive left eigenvector of A_+ associated with the reactive eigenvalue of A_+ . We label $L_4^-, L_5^\pm$ the eigenvectors of $\tilde{\mathbb{A}}_\pm$ associated with $-\mu_4^-$ and $-\mu_5^\pm$. By the block structure of $-\mu A_\pm + \mu^2 B_\pm + G_\pm$, spectral separation of μ_4^- and μ_5^- (and of μ_4^+ and μ_5^+), the eigenvectors $L_4^\pm$ and $L_5^\pm$ are analytical in λ , in a neighborhood of the origin (see for instance [Kat], II.1.4); in particular,

$$(3.27) \quad \underline{\ell}_4^+ = \ell_4^+ + O(\lambda), \quad \mu_4^+ b_+^{\text{tr}} \underline{\ell}_4^+ = O(\lambda).$$

The solutions of (3.21), seen as an equation in λ , are the eigenvalues of the matrix $-\mu \partial_v f_{|\pm}^\# + \mu^2 b_{1|\pm}^\#$. By (1.3) and the block structure of $A_\pm$, we find that the spectrum of $\partial_v f_{|\pm}^\#$ is

$$\sigma(\partial_v f_{|\pm}^\#) = \{-s(\varepsilon) - \sigma_\pm, \quad -s(\varepsilon), \quad -s(\varepsilon) + \sigma_\pm\}.$$

The eigenvalues of $\partial_v f_{|\pm}^\sharp$ are distinct, hence, by Rouché's theorem, the eigenvalues of $-\partial_v f_{|\pm}^\sharp + \mu b_{1|\pm}^\sharp$ are analytic in μ , for small μ , with expansions

$$(3.28) \quad \begin{aligned} \lambda_1 &= s + \sigma_\pm + \beta_1^\pm \mu + O(\mu^2), \\ \lambda_2 &= s + \beta_2^\pm \mu + O(\mu^2), \\ \lambda_3 &= s - \sigma_\pm + \beta_3^\pm \mu + O(\mu^2). \end{aligned}$$

By (H3) (Section 1.5), $\beta_j^\pm > 0$ for all j . Inversion of these expansions yields analytic functions $\mu_j^\pm$, called fluid modes, and defined in a neighborhood of the origin in $\mathbb{C}_\lambda$:

$$(3.29) \quad \begin{aligned} \mu_1^\pm &:= (s + \sigma_\pm)^{-1} \lambda - (s + \sigma_\pm)^{-3} \beta_1^\pm \lambda^2 + O(\lambda^3), \\ \mu_2^\pm &:= s^{-1} \lambda - s^{-3} \beta_2^\pm \lambda^2 + O(\lambda^3), \\ \mu_3^\pm &:= (s - \sigma_\pm)^{-1} \lambda - (s - \sigma_\pm)^{-3} \beta_3^\pm \lambda^2 + O(\lambda^3). \end{aligned}$$

By (3.18), the eigenvectors of $\tilde{\mathbb{A}}$ that are associated with these eigenvalues are

$$(3.30) \quad L_j^\pm(\lambda) = \begin{pmatrix} \underline{\ell}_j^\pm \\ \mu_j^\pm b_{\pm}^{\text{tr}} \underline{\ell}_j^\pm \end{pmatrix} \in \mathbb{C}^4 \times \mathbb{C}^3, \quad \underline{\ell}_j^\pm(0) = \underline{\ell}_j^\pm, \quad 1 \leq j \leq 3,$$

where the vectors $\underline{\ell}_1^\pm$, $\underline{\ell}_2^\pm$ and $\underline{\ell}_3^\pm$ are the left eigenvectors of $A_\pm$ associated with the fluid eigenvalues $-s - \sigma_\pm$, $-s$, and $-s + \sigma_\pm$; they have the form

$$(3.31) \quad \underline{\ell}_j^\pm := \begin{pmatrix} * & * & * & 0 \end{pmatrix}^{\text{tr}}, \quad 1 \leq j \leq 3.$$

The eigenvalues of $-\partial_v f_{|\pm}^\sharp + \mu b_{1|\pm}^\sharp$ being distinct, the associated eigenvectors are analytical as well, so that the $L_j^\pm$, $1 \leq j \leq 3$, are analytical in λ ; in particular,

$$(3.32) \quad \underline{\ell}_j^\pm = \underline{\ell}_j^\pm + O(\lambda), \quad \mu_j^\pm b_{\pm}^{\text{tr}} \underline{\ell}_j^\pm = O(\lambda).$$

Finally, the equation $\det(-\mu \partial_v f_{|\pm}^\sharp + \mu^2 b_{1|\pm}^\sharp) = 0$ has two non-zero solutions $\gamma_6^\pm, \gamma_7^\pm$, corresponding to the remaining two (fast) modes, solutions of

$$(3.33) \quad \kappa \tau_\pm^{-2} c^{-1} s \nu \mu^2 + (\kappa c^{-1} (s^2 - \Gamma \tau_\pm^{-2} e_\pm) + \nu s^2) \tau_\pm^{-1} \mu + s(s^2 - \sigma_\pm^2) = 0.$$

The Lax condition (1.9) implies that solutions of (3.33) are distinct and have small frequency expansions

$$(3.34) \quad \begin{aligned} \mu_6^\pm &= \gamma_6^\pm + O(\lambda), & \gamma_6^\pm &< 0, \\ \mu_7^\pm &= \gamma_7^\pm + O(\lambda), & \gamma_7^- &> 0, \quad \gamma_7^+ < 0. \end{aligned}$$

We label $L_6^\pm$ and $L_7^\pm$ the eigenvectors of $\tilde{\mathbb{A}}$ associated with $-\mu_6^\pm$ and $-\mu_7^\pm$. Again, by spectral separation, $L_6^\pm$ and $L_7^\pm$ are analytical in λ .

Lemma 3.5. *For some $r > 0$, equations (3.13) have analytical bases of solutions in $B(0, r)$,*

$$(3.35) \quad \tilde{\mathcal{B}}^\pm := \{\tilde{V}_j^\pm\}_{1 \leq j \leq 7}, \quad \tilde{V}_j^\pm := e^{-y\mu_j^\pm(\lambda)} L_j^\pm(\lambda),$$

where the eigenvalues $\mu_j^\pm$ are given in (3.22), (3.23), (3.29), and (3.34) and the eigenvectors associated with the slow modes are given in (3.25), (3.27), (3.30) and (3.32).

Proof. The above discussion describes analytical families $\mu_j^\pm$, $L_j^\pm$, such that the vectors $V_j^\pm$ defined in (3.35) are analytical solutions of (3.13). For $\lambda \neq 0$, the eigenvalues $\mu_j^\pm$ are simple, so that the families $\tilde{\mathcal{B}}^\pm$ define bases of equations (3.13). By inspection of the expansions at $\lambda = 0$, the families $\tilde{\mathcal{B}}^\pm$ define bases of equations (3.13) at $\lambda = 0$ as well. $\square$

The above low-frequency expansions of the eigenvalues show that

- (i) Equation $\tilde{W}' = \tilde{\mathbb{A}}_-(\lambda)\tilde{W}$ has a 3-dimensional subspace of solutions associated with slow modes (μ_j^- , $j = 1, 2, 3$) and 4-dimensional subspace of solutions associated with fast modes ($\mu_4^-, \mu_5^-, \mu_6^-, \mu_7^-$);
- (ii) Equation $\tilde{W}' = \tilde{\mathbb{A}}_+(\lambda)\tilde{W}$ has a 4-dimensional subspace of solutions associated with slow modes (μ_j^+ , $j = 1, 2, 3$, and μ_4^+) and a 3-dimensional subspace of solutions associated with fast modes ($\mu_5^+, \mu_6^+, \mu_7^+$).

3.1.3 Description of the essential spectrum

We adopt Henry's definition of the essential spectrum [He]:

Definition 3.6 (Essential spectrum). *Let $\mathcal{B}$ be a Banach space and $T : D(T) \subset \mathcal{B} \rightarrow \mathcal{B}$ a closed, densely defined operator. The essential spectrum of T , denoted by $\sigma_{ess}(T)$, is defined as the complement of the set of all λ such that λ is either in the resolvent set of T , or is an eigenvalue with finite multiplicity that is isolated in the spectrum of T .*

By Lemma 3.3, the matrix $\mathbb{A}_\pm(\lambda)$ has a non trivial center subspace if and only if $\lambda \in \mathcal{C}_\pm$,

$$\mathcal{C}_\pm := \{\lambda \in \mathbb{C}, \quad \det(-i\xi A_\pm - \xi^2 B_\pm + G_\pm - \lambda) = 0, \quad \text{for some } \xi \in \mathbb{R}\}.$$

The following Lemma can be found in [He] (Theorem A.2, Chapter 5 of [He], based on Theorem 5.1, Chapter 1 of [GK]):

Lemma 3.7. *The connected component of $\mathbb{C} \setminus (\mathcal{C}_- \cup \mathcal{C}_+)$ containing real $+\infty$ is a connected component of the complement of the essential spectrum of $L(\varepsilon)$.*




Figure 3: Spectrum of $L(\varepsilon)$.

The reactive eigenvalues of $-i\xi A_{\pm} - \xi^2 B_{\pm} + G_{\pm}$ are

$$\lambda = i\xi s - \xi^2 d - k\phi_{\pm}.$$

For small $|\xi|$, the fluid eigenvalues satisfy

$$\lambda = i\underline{\alpha}\xi - \beta\xi^2 + O(\xi^3), \quad \underline{\alpha} \in \mathbb{R}, \quad \beta > 0,$$

as described in Section 3.1.2; for large $|\xi|$, they satisfy

$$(3.36) \quad \lambda = -\xi^2(\bar{\alpha} + O(\xi^{-1})) \quad (\text{parabolic eigenvalues}),$$

with $\bar{\alpha} \in \{\nu\tau_{\pm}^{-1}, \kappa c^{-1}\tau_{\pm}^{-1}\}$, or

$$(3.37) \quad \lambda = is\xi + O(1) \quad (\text{hyperbolic eigenvalue}).$$

This implies that the essential spectrum is confined to the shaded area in Figure 3, the boundary of which is the union of an arc of parabola and two half-lines. (The origin $\lambda = 0$ is an eigenvalue, associated with eigenfunction $(\bar{U}^{\varepsilon})'$; the existence of bifurcation eigenvalues $\gamma(\varepsilon) \pm i\tau(\varepsilon)$ is assumed in Theorem 1.14, the proof of which is given in Section 5.)

Remark 3.8. *The essential spectrum, as given by Definition 3.6, is not stable under relatively compact perturbations (see [EE], Chapter 4, Example 2.2); namely, a domain of the complement of the essential spectrum of a (closed, densely defined) operator T is either a subset of the complement of the essential spectrum of $T + S$, or is filled with point spectrum of $T + S$, where S is a relatively compact perturbation of T .*

Remark 3.9. By the Fréchet-Kolmogorov theorem, L is a relatively compact perturbation of $L_{\pm}$. (This observation is the first step of the proof of Lemma 3.7, see Henry [He].) The pathology described in Remark 3.8 does not occur in the right half-plane here, as we know by an energy estimate that if λ is large and real, $\lambda \notin \sigma_p(L)$.

3.1.4 Gap Lemma and dual basis

Let Λ be the connected component of $\mathbb{C} \setminus (\mathcal{C}_- \cup \mathcal{C}_+)$ containing $\text{real } +\infty$.

Definition 3.10 (Stable and unstable subspaces at $\pm\infty$). Given $\lambda \in \Lambda \cup B(0, r)$, r as in Lemma 3.5, denote by $S(\tilde{\mathbb{A}}_{\pm}(\lambda))$ the stable subspace of $\tilde{\mathbb{A}}_{\pm}(\lambda)$ (i.e., the subspace of generalized eigenvectors associated with eigenvalues with negative real parts) and by $U(\tilde{\mathbb{A}}_{\pm}(\lambda))$ the unstable subspace of $\tilde{\mathbb{A}}_{\pm}$ (i.e., the subspace of generalized eigenvectors associated with eigenvalues with positive real parts). We define similarly $S(\mathbb{A}_{\pm}(\lambda))$ and $U(\mathbb{A}_{\pm}(\lambda))$.

By definition of $\mathcal{C}_{\pm}$, given $\lambda \in \Lambda$, the matrices $\mathbb{A}_{\pm}(\lambda)$ do not have purely imaginary eigenvalues, so that $S(\mathbb{A}_{\pm}(\lambda)) \oplus U(\mathbb{A}_{\pm}(\lambda)) = \mathbb{C}^7$, and $S(\tilde{\mathbb{A}}_{\pm}(\lambda)) \oplus U(\tilde{\mathbb{A}}_{\pm}(\lambda)) = \mathbb{C}^7$, for all $\lambda \in \Lambda$.

Lemma 3.11. The vector spaces $S(\tilde{\mathbb{A}}_{\pm}(\lambda))$ and $U(\tilde{\mathbb{A}}_{\pm}(\lambda))$ have analytical bases in Λ .

Proof. By simple-connectedness of Λ , the Lemma follows from a result of Kato ([Kat], II.4), that uses spectral separation in Λ . $\square$

Corollary 3.12. Equations (3.13) have analytical bases of solutions in Λ .

Proof. Basis elements of the stable and unstable spaces defined in Definition 3.10 are associated, through the flow of (3.13), with bases of solutions of (3.13). The matrices $\tilde{\mathbb{A}}_{\pm}$ depending analytically on λ , the flow of (3.13) is analytical in λ . $\square$

Lemma 3.13. For λ real and large, $\dim S(\mathbb{A}_+(\lambda)) = \dim S(\mathbb{A}_-(\lambda)) = 3$.

Proof. From Lemma 3.3, μ is an eigenvalue of $\mathbb{A}_{\pm}(\lambda)$ if and only if λ is an eigenvalue of $-\mu A_{\pm} + \mu^2 B_{\pm} + G_{\pm}$. As in Section 3.1.3, for large μ , the eigenvalues of $-\mu A_{\pm} + \mu^2 B_{\pm} + G_{\pm}$ are $s\mu + O(1)$ (hyperbolic mode) and $\nu\tau_{\pm}^{-1}\mu^2 + O(\mu)$, $\kappa\tau_{\pm}^{-1}c^{-1}\mu^2 + O(\mu)$, $d\mu^2 + O(\mu)$ (parabolic modes), c^{-1} as in Assumption 1.1. Inversion of these expansions gives three stable eigenvalues for both $\mathbb{A}_-$ and $\mathbb{A}_+$. $\square$

Remark 3.14. The above Lemma implies in particular that Λ is a domain of consistent splitting, as defined in [AGJ]. (See also Section 3.1 of [LyZ2].)

Given $\lambda \in \Lambda$, the flow of (3.13) associates basis elements of $S(\tilde{\mathbb{A}}_+(\lambda))$ with solutions of (3.13) which are exponentially decaying as $t \rightarrow +\infty$, and basis elements of $U(\tilde{\mathbb{A}}_-(\lambda))$ with solutions which are exponentially decaying as $t \rightarrow -\infty$. Similarly, the spaces $S(\tilde{\mathbb{A}}_-(\lambda))$ and $U(\tilde{\mathbb{A}}_+(\lambda))$ are associated with exponentially growing solutions, at $-\infty$ and $+\infty$ respectively.




Figure 4: Normal modes on the $-\infty$ side.

Definition 3.15 (Decaying and growing normal modes). *We call decaying dual normal mode at $\pm\infty$ any continuous family of dual normal modes $\{\lambda, \tilde{U}(\lambda)\}$, $\lambda \in B(0, r)$, r as in Lemma 3.5, such that for all $\lambda \in \Lambda \cap B(0, r)$, $\tilde{U}(\lambda)$ corresponds to a decaying solution of (3.13) at $\pm\infty$. Families of normal modes which are not decaying are growing. We define similarly decaying dual normal modes and growing dual normal modes.*

By continuity of the eigenvalues and spectral separation in Λ , if for some $\lambda \in \Lambda$ a continuous family of normal modes corresponds to a decaying (resp. growing) solution, then it corresponds for all $\lambda \in \Lambda$ to a decaying (resp. growing) solution.

By (1.9), (3.24) and (3.29), μ_1^+ , μ_2^+ , μ_3^+ and μ_4^+ are growing (in the sense of Definition 3.15) at $+\infty$, while μ_5^+ , μ_6^+ and μ_7^+ are decaying.

Similarly, μ_3^- , μ_5^- and μ_6^- are growing, while μ_1^- , μ_2^- , μ_4^- and μ_7^- are decaying.

The normal modes with which the characteristics of (1.1) are associated are pictured on Figures 5 and 4. In particular, slow normal modes associated with incoming characteristics are growing.

Definition 3.16 (Normal residuals). *A map $(y, \lambda) \rightarrow \Theta^+(y, \lambda) \in \mathbb{C}^7$ defined on $[y_0, +\infty) \times B(0, r)$, for some $y_0 > 0$, $r > 0$, is said to belong to the class of normal residuals if it satisfies the estimates*

$$|\Theta^+| \leq C, \quad |\partial_y \Theta^+| \leq C(|\lambda| + e^{-\theta|y|}).$$

for some $\theta > 0$ and $C > 0$, uniformly in $y \geq y_0$ and $\lambda \in B(0, r)$.

We define similarly the class of normal residuals on $(-\infty, -y_0) \times B(0, r)$.




Figure 5: Normal modes on the $+\infty$ side.

Lemma 3.17 (Fast dual modes). *Equations (3.8) has solutions*

$$\tilde{W}_4^-, \tilde{W}_5^+, \tilde{W}_6^+, \tilde{W}_7^+ \quad (\text{growing}) \quad \text{and} \quad \tilde{W}_5^-, \tilde{W}_6^-, \tilde{W}_7^- \quad (\text{decaying}),$$

which for $\lambda \in B(0, r)$, r possibly smaller than in Lemma 3.5, satisfy

$$(3.38) \quad \tilde{W}_j^\pm = e^{-y\mu_j^\pm(\lambda)} (L_j^\pm(0) + e^{-\theta|y|} \tilde{\Theta}_{1j}^\pm + \lambda \tilde{\Theta}_{2j}^\pm), \quad y \gtrless \pm y_0,$$

for some $y_0 > 0$ independent of λ , where the constant vectors $L_j^\pm(0)$ are defined in Section 3.1.2, and $\tilde{\Theta}_{1j}^\pm, \tilde{\Theta}_{2j}^\pm$ are normal residuals in the sense of Definition 3.16.

Proof. With the description of the normal modes in Lemma 3.5, this is a direct application of the Gap Lemma (for instance in the form of Proposition 9.1 of [MaZ3]). $\square$

Lemma 3.18 (Slow dual modes). *Equation (3.8) has solutions*

$$\tilde{W}_1^-, \tilde{W}_2^- \quad (\text{growing}) \quad \text{and} \quad \tilde{W}_3^-, \tilde{W}_1^+, \tilde{W}_2^+, \tilde{W}_3^+, \tilde{W}_4^+ \quad (\text{decaying}),$$

which for $\lambda \in B(0, r)$, r possibly smaller than in Lemma 3.5, satisfy

$$(3.39) \quad \tilde{W}_j^\pm = e^{-y\mu_j^\pm(\lambda)} (L_j^\pm(0) + \lambda \tilde{\Theta}_j^\pm), \quad y \gtrless \pm y_0,$$

for some $y_0 > 0$ independent of λ , where the constant vectors $L_j^\pm(0)$ are defined in Section 3.1.2, and $\tilde{\Theta}_j^\pm$ are normal residuals.

Proof. The Conjugation Lemma ([MeZ]; Lemma 3.1 of [MaZ3]) implies that there exists a family of matrix-valued applications $\{\tilde{\Theta}^+(\cdot, \lambda)\}_{\lambda \in B(0, r)}$, for some $r > 0$ possibly smaller than in Lemma 3.5, such that the matrix $\text{Id} + \tilde{\Theta}^+$ is invertible for all λ and y , the application $\tilde{\Theta}^+$ is smooth in y and analytical in λ , with exponential bounds

$$|\partial_\lambda^j \partial_x^k \tilde{\Theta}^+| \leq C_{jk} e^{-\theta y}, \quad \text{for some } \theta > 0, C_{jk} > 0, \text{ for } y \geq y_0,$$

for some $y_0 > 0$, and such that any solution $\tilde{W}$ of (3.8) has the form

$$(3.40) \quad \tilde{W} = (\text{Id} + \tilde{\Theta}^+) \tilde{V}^+, \quad \text{for } y \geq y_0,$$

where $\tilde{V}^+$ is a dual normal mode, and, conversely, if $\tilde{V}^+$ is a dual normal mode, then $\tilde{W}$ defined by (3.40) solves (3.8) on $y \geq y_0$.

Equation (3.8) at $\lambda = 0$ has a four-dimensional subspace of constant solutions; let $\{\tilde{W}_j^0\}_{1 \leq j \leq 4}$ be a generating family. The normal modes with which, through (3.40), the $\tilde{W}_j^0$ are associated are slow normal modes. Hence, by Lemma 3.5, there exist coordinates c_{jk} such that

$$\tilde{W}_j^0 = (\text{Id} + \tilde{\Theta}^+(\cdot, 0)) \sum_{1 \leq k \leq 4} c_{jk} L_k^+(0), \quad y \geq y_0,$$

which implies in particular that the matrix $c := (c_{jk})_{1 \leq j, k \leq 4}$ is invertible. Then, for $1 \leq j \leq 4$,

$$(\text{Id} + \tilde{\Theta}^+(\cdot, 0)) L_j^+(0) = \sum_{1 \leq k \leq 4} (c^{-1})_{jk} \tilde{W}_k^0,$$

in particular, $(\text{Id} + \tilde{\Theta}^+(\cdot, 0)) L_j^+(0)$ is constant, hence, by exponential decay of $\tilde{\Theta}^+$, equal to $L_j^+(0)$. We can conclude that, for $1 \leq j \leq 4$,

$$\tilde{W}_j^+ := (\text{Id} + \tilde{\Theta}^+) \tilde{V}_j^+$$

(where $\tilde{V}_j^+$ is defined in Lemma 3.5) is a solution of (3.8) on $y \geq y_0$, which can be put in the form (3.39).

The proof on the $-\infty$ side is based similarly on the decomposition of the fluid components of the $\tilde{W}_j^0$ onto the (fluid) dual slow modes $\tilde{V}_j^-$, for $1 \leq j \leq 3$. $\square$

3.1.5 Duality relation and forward basis

We use the duality relation, introduced in [MaZ3],

$$(3.41) \quad \tilde{W}^{\text{tr}} \mathcal{S} W = 1$$

that relates solutions W of the forward equation (3.3) with solutions $\tilde{W}$ of the adjoint equation (3.8) through the conjugation matrix in (τ, w, bw') coordinates

$$\mathcal{S} := \begin{pmatrix} -A_{11} & -A_{12} & 0 \\ -A_{21} & -A_{22} & \text{Id}_{\mathbb{C}^3} \\ 0 & -\text{Id}_{\mathbb{C}^3} & 0 \end{pmatrix},$$

where A is the convection matrix defined in (1.16). Namely, W is a solution of (3.3) if and only if it satisfies (3.41) for all solution $\tilde{W}$ of (3.8), and conversely $\tilde{W}$ is a solution of (3.8) if and only if it satisfies (3.41) for all solution W of (3.3). (See Lemma 4.2, [MaZ3]; note that the reactive term contains no derivative, hence does not play any role here.)

Remark that there exist vectors $r_k^\pm$ such that

$$(3.42) \quad \ell_j^\pm A_\pm r_k^\pm = -\delta_{jk}, \quad 1 \leq j, k \leq 4.$$

Let $R_k^\pm$ be vectors of the form

$$(3.43) \quad R_k^\pm := \begin{pmatrix} r_k^\pm \\ * \end{pmatrix} + e^{-\theta} \Theta_{1k}^\pm,$$

where for $1 \leq k \leq 4$, $r_k^\pm$ are given by (3.43), and where $\Theta_{1k}^\pm$ are normal residuals. With the notation of Lemmas 3.17 and 3.18, let

$$(3.44) \quad \bar{L}_j^\pm := \begin{cases} L_j^\pm(0) & \text{if } \mu_j^\pm \text{ is slow,} \\ L_j^\pm(0) + e^{-\theta|y|} \bar{\Theta}_{1j}^\pm & \text{if } \mu_j^\pm \text{ is fast.} \end{cases}$$

Lemma 3.19 (Forward and dual basis). *For some $r > 0$ and $y_0 > 0$,*

- *equation (3.3) has analytical bases of solutions $\{W_1^\pm, \dots, W_7^\pm\}_{\lambda \in \Lambda \cup B(0, r)}$, for $y \gtrless \pm y_0$;*
- *equation (3.8) has analytical bases of solutions $\{\bar{W}_1^\pm, \dots, \bar{W}_7^\pm\}_{\lambda \in \Lambda \cup B(0, r)}$, for $y \gtrless \pm y_0$,*

such that for $\lambda \in B(0, r)$,

$$(3.45) \quad W_j^\pm = e^{x\mu_j^\pm(\lambda)} (R_j^\pm + \lambda \Theta_j^\pm), \quad y \gtrless \pm y_0,$$

$$(3.46) \quad \bar{W}_j^\pm = e^{-y\mu_j^\pm(\lambda)} (\bar{L}_j^\pm + \lambda \bar{\Theta}_j^\pm), \quad y \gtrless \pm y_0,$$

where $R_j^\pm$ and $\bar{L}_j^\pm$ are defined in (3.43) and (3.44), and $\Theta_j^\pm$ and $\bar{\Theta}_j^\pm$ are normal residuals; the fast forward modes W_4^- and W_7^+ satisfy also

$$(3.47) \quad W_j^\pm(x, \lambda) = \begin{pmatrix} (\bar{U}^\varepsilon)'(x) \\ * \end{pmatrix} + \lambda \underline{\Theta}_j^\pm(x, \lambda), \quad x \gtrless \pm y_0,$$

where $|\underline{\Theta}_j^\pm| + |\partial_x \underline{\Theta}_j^\pm| \leq C e^{-\theta|x|}$, for some $C, \theta > 0$, uniformly in $\lambda \in B(0, r)$.

Proof. Given a family $\{F_1, \dots, F_7\}$ of vectors in $\mathbb{C}^7$, let $\text{col}(F_j)$ denote the 7×7 matrix $\text{col}(F_j) := \begin{pmatrix} F_1 & \dots & F_7 \end{pmatrix}$.

Let y_0, r , and $\tilde{W}_j^\pm$ as in Lemma 3.17 and 3.18. For all $\lambda \in \Lambda \cup B(0, r)$, the families $\{\tilde{W}_1^-, \dots, \tilde{W}_n^-\}$ and $\{\tilde{W}_1^+, \dots, \tilde{W}_n^+\}$ are bases of solutions of (3.8), on $y \leq -y_0$ and $y \geq y_0$ respectively. In particular, the 7×7 matrices $\tilde{\mathbf{W}}^{0\pm} := \text{col}(\tilde{W}_j^\pm)$ are invertible for all $\lambda \in B(0, r)$ and $y \gtrless \pm y_0$. Let

$$(3.48) \quad \mathbf{W}^{0\pm} := ((\tilde{\mathbf{W}}^{0\pm})^{\text{tr}} \mathcal{S})^{-1} =: \text{col}(W_k^{0\pm}).$$

For the forward modes $W_j^{0\pm}$ defined in (3.48) to satisfy the low-frequency description

$$(3.49) \quad W_j^{0\pm} = e^{x\mu_j^\pm(\lambda)} (R_j^{0\pm} + e^{-\theta|x|} \Theta_{1j}^{0\pm} + \lambda \Theta_{2j}^{0\pm}), \quad y \gtrless \pm y_0,$$

where $R_j^{0\pm}$ are constant vectors and $\Theta_{\star j}^{0\pm}$ are normal residuals, it suffices, by (3.41), that the matrices $\mathbf{R}^{0\pm} := \text{col}(R_j^{0\pm})$ and $\Theta_\star^{0\pm} := \text{col}(\Theta_{\star j}^{0\pm})$ satisfy

$$(3.50) \quad \mathbf{L}^{\text{tr}} \mathbf{S} \mathbf{R}^0 = \text{Id}_{\mathbb{C}^7},$$

$$(3.51) \quad (\mathbf{L} + e^{-\theta|x|} \tilde{\Theta}_1)^{\text{tr}} \mathbf{S} \Theta_1^0 = -\tilde{\Theta}_1^{\text{tr}} \mathbf{S} \mathbf{R}^0,$$

$$(3.52) \quad (\mathbf{L} + e^{-\theta|x|} \tilde{\Theta}_1^0 + \lambda \tilde{\Theta}_2)^{\text{tr}} \mathbf{S} \Theta_2^0 = -\tilde{\Theta}_2^{\text{tr}} \mathbf{S} \mathbf{R}^0,$$

where $\mathbf{L}^\pm := \text{col}(L_j^\pm(0))$ and $\tilde{\Theta}_\star^\pm := \text{col}(\tilde{\Theta}_{\star j}^\pm)$ appear in the low-frequency description of the $\tilde{W}_j^\pm$. In (3.50)-(3.52), the $\pm$ exponents are omitted. The matrices $\mathbf{L}^\pm$ being invertible, (3.50) (with $+$ or $-$) has a unique solution, and, for y_0 large enough and r small enough, equations (3.51) and (3.52) have unique solutions in the class of normal residuals. Note that for $1 \leq j, k \leq 4$, equation (3.50) reduces to (3.42), up to exponentially decaying terms, so that the vectors $R_k^{0\pm}$ have the form (3.43).

Remark now that $(\bar{U}^\varepsilon)'$ satisfies $L(\varepsilon)(\bar{U}^\varepsilon)' = 0$, and decays at both $-\infty$ and $+\infty$, hence $(\bar{U}^\varepsilon)'$ is associated with decaying fast normal modes; by Lemma 3.17, there exist constants $c_j^\pm$, such that

$$(3.53) \quad \begin{pmatrix} (\bar{U}^\varepsilon)'(y) \\ * \end{pmatrix} = c_4^- W_4^{0-}|_{\lambda=0} = \sum_{5 \leq j \leq 7} c_j^+ W_j^{0+}|_{\lambda=0}.$$

We may assume, without loss of generality, that $c_7^+ \neq 0$. Let now

$$\begin{aligned} \mathbf{W}^- &:= \begin{pmatrix} W_1^{0-} & W_2^{0-} & W_3^{0-} & c_4^- W_4^{0-} & W_5^{0-} & W_6^{0-} & W_7^{0-} \end{pmatrix}, \\ \mathbf{W}^+ &:= \begin{pmatrix} W_1^{0+} & W_2^{0+} & W_3^{0+} & W_4^{0+} & W_5^{0+} & W_6^{0+} & \sum_{j=5}^7 c_j^+ W_j^{0+} \end{pmatrix}, \end{aligned}$$

and $\mathbf{W}^\pm =: \text{col}(W_j^\pm)$. These forward modes satisfy (3.45) and (3.47). Let finally $\tilde{\mathbf{W}}^{\pm \text{tr}} := (\mathbf{S} \mathbf{W}^\pm)^{-1} =: \text{col}(\tilde{W}_j^\pm)$, so that, in particular, the slow modes of $\tilde{\mathbf{W}}^{0\pm}$ and $\tilde{\mathbf{W}}^\pm$ coincide. We can prove as above that the low-frequency description (3.49) of the forward modes carries over to the dual modes through the duality relation, so that (3.46) is satisfied. $\square$

3.1.6 The resolvent kernel

Let

$$L^2(\Omega, \mathcal{D}'(\mathbb{R})) := \{\phi \in \mathcal{D}'(\Omega \times \mathbb{R}), \text{ for all } \varphi \in \mathcal{D}(\mathbb{R}), \langle \phi, \varphi \rangle \in L^2(\Omega)\}$$

A linear continuous operator $T : L^2(\mathbb{R}) \rightarrow L^2(\mathbb{R})$ operates on $L^2(\mathbb{R}, \mathcal{D}'(\mathbb{R}))$, by $\langle T\phi, \varphi \rangle := T\langle \phi, \varphi \rangle$. Let $\tau_{(\cdot)}\delta \in L^2(\mathbb{R}, \mathcal{D}'(\mathbb{R}))$ be defined by $\langle \tau_x\delta, \varphi \rangle = \varphi(x)$, for all $x \in \mathbb{R}$.

Definition 3.20 (Resolvent kernel). *Given λ in the resolvent set of $L(\varepsilon)$, define the resolvent kernel $\mathcal{G}_\lambda$ of $L(\varepsilon)$ as an element of $L^2(\mathbb{R}_x, \mathcal{D}'(\mathbb{R}_y))$ by*

$$\mathcal{G}_\lambda := (L(\varepsilon) - \lambda)^{-1} \tau_{(\cdot)}\delta.$$

Given $y \in \mathbb{R}$, let $s_y = \text{sgn}(y)$, and

$$\tilde{\mathbf{D}}(y) := \{j, \quad \tilde{\mu}_j^{s_y} \text{ slow and decaying}\},$$

so that

$$\tilde{\mathbf{D}}(y) = \{3\}, \quad \text{if } y < 0, \quad \tilde{\mathbf{D}}(y) = \{1, 2, 3, 4\}, \quad \text{if } y > 0.$$

Given $x, y \in \mathbb{R}$, let $\mathbf{D}(x, y)$ be the set of all (j, k) such that for all x, y , for $\Re\lambda > 0$ and $|\lambda|$ small enough, $\Re(\mu_j^{s_x}x - \mu_k^{s_y}y) < 0$, that is,

$$\begin{aligned} \mathbf{D}(x, y) = & \{(j, k), \quad \mu_j^{s_x} \text{ and } \tilde{\mu}_k^{s_y} \text{ slow and decaying}\} \\ & \bigcup \{(j, j), \quad s_x = s_y, \quad |y| < |x|, \quad \mu_j^{s_x} \text{ slow and decaying}\} \\ & \bigcup \{(j, j), \quad s_x = s_y, \quad |x| < |y|, \quad \tilde{\mu}_j^{s_y} \text{ slow and decaying}\}, \end{aligned}$$

so that

$$\mathbf{D}(x, y) := \begin{cases} \{(1, 1), (2, 2), (1, 3), (2, 3)\}, & x \leq y \leq 0, \\ \{(1, 3), (2, 3), (3, 3)\}, & y \leq x \leq 0, \\ \emptyset, & y \leq 0 \leq x, \\ \{(j, k), \quad 1 \leq j \leq 4, 1 \leq k \leq 2\}, & x \leq 0 \leq y, \\ \{(j, j), \quad 1 \leq j \leq 4\}, & 0 \leq x \leq y, \\ \emptyset, & 0 \leq y \leq x. \end{cases}$$

Define now the excited term

$$\mathcal{E}_\lambda(x, y) := \lambda^{-1} (\bar{U}^\varepsilon)'(x) \sum_{j \in \tilde{\mathbf{D}}(y)} [c_{j, s_y}^0] \ell_j^{s_y \mathbf{tr}} e^{-y \mu_j^{s_y}(\lambda)},$$

and the scattered term

$$\mathcal{S}_\lambda(x, y) := \sum_{(j, k) \in \mathbf{D}(x, y)} [c_{k, s_y}^{j, s_x}] r_j^{s_x} \ell_k^{s_y \mathbf{tr}} e^{x \mu_j^{s_x}(\lambda) - y \mu_k^{s_y}(\lambda)},$$

where the vectors $\ell_j^\pm$ are defined in (3.26) and (3.31), the vectors $r_j^\pm$ are defined in (3.42), and the transmission coefficients $[c_{k, \pm}^0]$ and $[c_{k, \pm}^{j, \pm}]$ are constants.

Proposition 3.21. *Under (1.18), for $\lambda \in B(0, r)$, the radius r being possibly smaller than in Lemma 3.5, there exist transmission coefficients $[c_{k,\pm}^0]$ and $[c_{k,\pm}^{j,\pm}]$ such that the resolvent kernel decomposes as*

$$\mathcal{G}_\lambda = \mathcal{E}_\lambda + \mathcal{S}_\lambda + \mathcal{R}_\lambda,$$

where $\mathcal{R}_\lambda$ satisfies

$$\begin{aligned} |\partial_x^\alpha \partial_y^{\alpha'} \mathcal{R}_\lambda| &\leq C e^{-\theta|x-y|} + C \lambda^{\alpha'} e^{-\theta|x|} \sum_{j \in \bar{\mathbf{D}}(y)} e^{-y \mu_j^{sy}} \\ &\quad + C (\lambda^{1+\min(\alpha, \alpha')} + \lambda^\alpha e^{-\theta|x|}) \sum_{(j,k) \in \mathbf{D}(x,y)} e^{x \mu_k^{sx} - y \mu_j^{sy}}, \end{aligned}$$

for $\alpha \in \{0, 1, 2, \dots\}$, $\alpha' \in \{0, 1\}$, for some $C, \theta > 0$, uniformly in x, y and $\lambda \in B(0, r)$.

Proof. The duality relation (3.41) allows to apply Proposition 4.6 of [MaZ3] (and its Corollary 4.7), which describes $\mathcal{G}_\lambda$ as sums of pairings of forward and dual modes, for λ in the intersection of Λ and the resolvent set of L . By Lemma 3.19, $\mathcal{G}_\lambda$ extends as a meromorphic map on $B(0, r)$.

The excited term $\mathcal{E}_\lambda$ comprises the pole terms, corresponding to pairings of a fast, decaying forward mode associated with the derivative of the background wave with a slow, decaying dual mode, i.e. $W_7^+/\bar{W}_3^-$ for $y \leq 0$ and $W_4^-/\bar{W}_j^+$ for $y \geq 0$, $1 \leq j \leq 4$.

The next-to-leading order term is the scattered term $\mathcal{S}_\lambda$. It corresponds to pairings of a slow forward mode with a slow dual mode. For $y \leq 0$, the scattered term comprises only fluid modes. For $y \leq 0 \leq x$ and for $0 \leq y \leq x$, the scattered term vanishes, as there are no outgoing modes to the right of the shock (see Figures 1 and 4).

By the Evans function condition (1.18) and Lemma 6.11 of [MaZ3], the residual $\mathcal{R}_\lambda$ does not contain any pole term; it comprises:

- (a) the contribution of the normal residuals to the fast forward/slow dual pairings involving the derivative of the background profile,
- (b) the fast forward/slow dual pairings not involving the derivative of the background profile,
- (c) the contribution of the normal residuals to the slow forward/slow dual pairings, and
- (d) the slow forward/fast dual pairings.

Term (a) is bounded by the first two terms in the upper bound for $\mathcal{R}_\lambda$. Term (b) is smaller than term (a) by a $O(\lambda)$ factor. Term (c) is bounded by the third term in the upper bound. By the Lax condition (1.9), the Evans function condition (1.18) and Lemma 6.11 of [MaZ3], term (d) is also bounded by the third term. $\square$

3.1.7 The Evans function

By Lemma 3.13, for all $\lambda \in \Lambda$, the dimensions of $U(\mathbb{A}_-(\lambda))$ and $S(\mathbb{A}_+(\lambda))$, the vector spaces associated with decaying solutions of (3.3) at $-\infty$ and $+\infty$, add up to the full dimension of the ambient space:

$$\dim U(\mathbb{A}_-(\lambda)) + \dim S(\mathbb{A}_+(\lambda)) = 7.$$

Definition 3.22 (Evans function). *On $\Lambda \cup B(0, r)$, define the Evans function as*

$$D(\varepsilon, \lambda) := \det(W_1^-, W_2^-, W_4^-, W_7^-, W_5^+, W_6^+, W_7^+)|_{x=0}.$$

The Evans function D satisfies Proposition 1.8; it has a zero at $\lambda = 0$, as reflected in equality (3.53).

3.2 Inverse Laplace transform

Similarly as in Section 3.1.6 (or Section 2 of [MaZ3]), define the Green function of $L(\varepsilon)$ as

$$(3.54) \quad \mathcal{G} := e^{tL(\varepsilon)} \tau_{(\cdot)} \delta,$$

where $\{e^{tL(\varepsilon)}\}_{t \geq 0}$ is the semi-group generated by $L(\varepsilon)$. That is, the kernel of the integral operator $e^{t_0 L(\varepsilon)}$ is the Green function $\mathcal{G}$ evaluated at $t = t_0$.

Assuming (1.18), the inverse Laplace transform representation of the semi-group by the resolvent operator (see for instance [Pa] Theorem 7.7; [Z3] Proposition 6.24) yields

$$(3.55) \quad \mathcal{G}(\varepsilon, x, t; y) = \frac{1}{2\pi i} \text{P.V.} \int_{\eta_0 - i\infty}^{\eta_0 + i\infty} e^{\lambda t} \mathcal{G}_\lambda(\varepsilon, x, y) d\lambda,$$

for $\eta_0 > 0$ sufficiently large.

3.2.1 Pointwise Green function bounds

Introduce the notations

$$\text{erfn}(y) := \int_{-\infty}^y e^{-z^2} dz,$$

and let, for $y < 0$:

$$(3.56) \quad \mathbf{e} := [c_{3,-}^0] \ell_3^{-\text{tr}} \left(\text{erfn} \left(\frac{y + a_3^- t}{\sqrt{4\beta_j^- t}} \right) - \text{erfn} \left(\frac{y - a_3^- t}{\sqrt{4\beta_j^- t}} \right) \right),$$

for $y > 0$:

$$(3.57) \quad \mathbf{e} := \sum_{1 \leq j \leq 4} [c_{j,+}^0] \ell_j^{+\text{tr}} \left(\text{erfn} \left(\frac{y + a_j^+ t}{\sqrt{4\beta_j^+ t}} \right) - \text{erfn} \left(\frac{y - a_j^+ t}{\sqrt{4\beta_j^+ t}} \right) \right),$$

and

$$(3.58) \quad \mathcal{E}(\varepsilon, x, t; y) := (\bar{U}^\varepsilon)'(x) \mathbf{e}(\varepsilon, t; y).$$

In (3.56)-(3.57) and below, the $\{a_j^\pm\}_{1 \leq j \leq 4}$ are the characteristic speeds, i.e. the limits at $\pm\infty$ of the eigenvalues of $\partial_U F(\varepsilon, \bar{U}^\varepsilon)$, ordered as in (1.3), the $\beta_j^\pm$, $1 \leq j \leq 3$, are the positive diffusion rates that were introduced in (3.28) (and which depend on ε , as do the characteristic speeds), and $\beta_4^+ := d$, the species diffusion coefficient.

Let for $y < 0$:

$$(3.59) \quad \begin{aligned} \mathcal{S} := & \chi_{\{t \geq 1\}} \sum_{1 \leq j \leq 2} r_j^- \ell_j^- \text{tr} (4\pi \beta_j^- t)^{-\frac{1}{2}} e^{-(x-y-a_j^- t)^2 / 4\beta_j^- t} \\ & + \chi_{\{t \geq 1\}} \frac{e^{-x}}{e^{-x} + e^x} \left(r_3^- \ell_3^- \text{tr} (4\pi \beta_3^- t)^{-\frac{1}{2}} e^{-(x-y+(s-\sigma_-)t)^2 / 4\beta_3^- t} \right. \\ & \quad \left. + \sum_{1 \leq j \leq 2} [c_{3,-}^{j,-}] r_j^- \ell_3^- \text{tr} (4\pi \beta_{3,-}^{j,-} t)^{-\frac{1}{2}} e^{-(x-z_{3,-}^{j,-})^2 / 4\beta_{3,-}^{j,-} t} \right), \end{aligned}$$

and for $y > 0$:

$$(3.60) \quad \begin{aligned} \mathcal{S} := & \chi_{\{t \geq 1\}} \frac{e^x}{e^{-x} + e^x} \sum_{1 \leq j \leq 4} r_j^+ \ell_j^+ \text{tr} (4\pi \beta_j^+ t)^{-\frac{1}{2}} e^{-(x-y-a_j^+ t)^2 / 4\beta_j^+ t} \\ & + \chi_{\{t \geq 1\}} \frac{e^{-x}}{e^{-x} + e^x} \sum_{1 \leq j \leq 4} \sum_{1 \leq k \leq 2} [c_{j,+}^{k,-}] r_k^- \ell_j^+ \text{tr} (4\pi \beta_{j,+}^{k,-} t)^{-\frac{1}{2}} e^{-(x-z_{j,+}^{k,-})^2 / 4\beta_{j,+}^{k,-} t}, \end{aligned}$$

where the indicator function $\chi_{\{t \geq 1\}}$ is identically equal to 1 for $t \geq 1$ and 0 otherwise, and

$$z_{j,\pm}^{k,\pm} := a_j^\pm (t - |y| |a_k^\pm|^{-1}), \quad \beta_{k,\pm}^{j,\pm} := \frac{|x|}{|a_j^\pm| t} \beta_j^\pm + \frac{|y|}{|a_k^\pm| t} \left(\frac{a_j^\pm}{a_k^\pm} \right)^2 \beta_k^\pm.$$

Let

$$(3.61) \quad \mathcal{H} := h(\varepsilon, t, x, y) \tau_{x+st} \delta, \quad h \ell_4^+ \equiv 0,$$

where the notation $\tau_{(\cdot)} \delta$ was introduced at the beginning of Section 3.1.6.

Let finally $\mathcal{S}_0$ be the scattered term defined in (3.59)-(3.60) in which $[c_{k,\pm}^{j,\pm}] = 1$ for all j, k .

Proposition 3.23. *Under (1.18), there exists transmission coefficients $[c_{j,\pm}^0]$ and $[c_{k,\pm}^{j,\pm}]$, satisfying*

$$(3.62) \quad \begin{cases} [c_{4+}^0] = 0, \\ r_k^* = [c_{k,*}^0] (v_+^\varepsilon - v_-^\varepsilon) + [c_{k,*}^{1,-}] r_1^- + [c_{k,*}^{2,-}] r_2^-, \\ [c_{4,+}^{1,-}] = [c_{4,+}^{2,-}] = 0, \end{cases}$$

where $1 \leq k \leq 3$ if $\star = +$ and $1 \leq k \leq 2$ if $\star = -$, $U_{\pm}^{\varepsilon} =: (v_{\pm}^{\varepsilon}, z_{\pm}^{\varepsilon})$, such that the Green function $\mathcal{G}(\varepsilon, x, t; y)$ defined in (3.54) may be decomposed as a sum of hyperbolic, excited, scattered, and residual terms, as follows:

$$(3.63) \quad \mathcal{G} = \mathcal{H} + \mathcal{E} + \mathcal{S} + \mathcal{R},$$

where $\mathcal{H}$, $\mathcal{E}$ and $\mathcal{S}$ are defined in (3.56)-(3.61), with the estimates

$$(3.64) \quad \begin{aligned} |\partial_t^k \partial_x^\alpha \partial_y^{\alpha'} h| &\leq C e^{-\theta t}, \\ |\partial_x^\alpha \partial_y^{\alpha'} \mathcal{R}| &\leq C e^{-\theta(|x-y|+t)} \\ &+ C \left(t^{-\frac{1}{2}(1+\alpha+\alpha')} (t+1)^{-\frac{1}{2}} + e^{-\theta t} \right) e^{-(x-y)^2/Mt} \\ &+ C \left((t^{-\frac{1}{2}} + e^{-\theta|x|}) t^{-\frac{1}{2}(\alpha+\alpha')} + \alpha' t^{-\frac{1}{2}} e^{-\theta|y|} \right) |\mathcal{S}_0|. \end{aligned}$$

uniformly in ε , for $k \in \{0, 1\}$, $\alpha \in \{0, 1, 2\}$, $\alpha' \in \{0, 1\}$, for some $\theta, C, M > 0$.

Proof. We only check (3.62), as decomposition (3.63) and bounds (3.64) are easily deduced from Proposition 7.1 of [MaZ3] and Proposition 7.3 of [LRTZ]. (See also Proposition 3.7 of [TZ2], especially equations (3.30)-(3.33) and (3.38).)

The description of the residue of $\mathcal{G}_\lambda$ at $\lambda = 0$ for $y < 0$ and $y > 0$ implies

$$[c_{3,-}^0] \ell_3^- = [c_{4,+}^0] \ell_4^+ + \sum_{1 \leq j \leq 3} [c_{j,+}^0] \ell_j^+,$$

corresponding to equation (1.34) in [MaZ3]. The (reactive) left eigenvector vector ℓ_4^+ being orthogonal to the (fluid) left eigenspace $\text{span}\{\ell_j^\pm\}_{1 \leq j \leq 3}$ (see (3.26) and (3.31)), this implies (3.62)(i).

Given $U_0 \in L^1$, the estimates for $\mathcal{H}$ and $\mathcal{R}$ imply

$$\lim_{t \rightarrow +\infty} \int_{\mathbb{R}^2} (\mathcal{H} + \mathcal{R}) U_0 dy dx = 0.$$

Hence, by conservation of mass in the fluid variables, (1.5), for all $U_0 \in L^1$,

$$(3.65) \quad \lim_{t \rightarrow +\infty} \int_{\mathbb{R}^2} \pi(\mathcal{E} + \mathcal{S}) U_0 dy dx = \int_{\mathbb{R}} \pi U_0 dy.$$

where $\pi : \mathbb{C}_{v,z}^4 \rightarrow \mathbb{C}_v^3$ is defined by $\pi(v, z) := v$. (Equation (3.65) corresponds to (1.33) and (7.60) in [MaZ3].) Taking $U_0 \in \text{span}\{\ell_j^\pm\}_{1 \leq j \leq 3}$, we find (3.62)(ii), and taking U_0 parallel to ℓ_4^+ , we find (3.62)(iii). $\square$

Remark 3.24. The terms $\mathcal{E}$ and $\mathcal{S}$ correspond to the low-frequency part of representation of $\mathcal{G}$ by inverse Laplace transform of the resolvent kernel $\mathcal{G}_\lambda$, while the term $\mathcal{H}$ corresponds to the high-frequency part. As observed in [MaZ3], for low frequencies, the resolvent kernel in the case of real (physical) viscosity obeys essentially

the same description as in the artificial (Laplacian) viscosity case, hence the estimates on $\mathcal{E}$ and $\mathcal{S}$ follow by the analysis in [LRTZ] of the corresponding artificial viscosity system, specialized to the case of strong detonations (more general waves were treated in [LRTZ]). The estimate of the terms $\mathcal{H}$ and $\mathcal{R}$ follows exactly as for the nonreactive case treated in [MaZ3, Z2].

Remark 3.25. Bound (3.64)(ii) is implied by bounds (7.1)-(7.4) of Proposition 7.1 of [MaZ3] and bounds (3.30), (3.32) and (3.38) of Proposition 3.7 of [TZ2].

Here the contribution of the hyperbolic, delta-function terms to the upper bounds for the spatial derivatives of $\mathcal{R}$ is absorbed in $\mathcal{H}$, and the short-time, $t \leq |a_k^\pm||y|$, contributions of the scattered terms are absorbed in the generic parabolic residual term $e^{-\theta t}e^{-(x-y)^2/Mt}$.

Corollary 3.26. The excited terms $\mathcal{E}_\lambda$ and $\mathcal{E}$ contain only fluid terms: $\mathcal{E}_\lambda \ell_4^+ \equiv 0$ and $\mathcal{E} \ell_4^+ \equiv 0$.

Proof. The equality $\mathcal{E} \ell_4^+ \equiv 0$ follows from (3.62)(i). The resolvent kernel $\mathcal{G}_\lambda$ is the Laplace transform of the Green function $\mathcal{G}$, so that the coefficients $[c_{j,\pm}^0]$ in Propositions 3.21 and 3.23 must agree. Hence, (3.62)(i) implies also $\mathcal{E}_\lambda \ell_4^+ \equiv 0$. $\square$

Corollary 3.27. For all $\eta > 0$, for some $C, M > 0$, some $\theta_1(\eta, s) > 0$, the following bounds hold, for $\alpha \in \{0, 1, 2\}$:

$$(3.66) \quad \begin{aligned} |e^{-\eta y^+} \partial_x^\alpha \mathcal{S} \ell_4^+| &\leq C e^{-\theta_1 t} e^{-\eta|x-y|/2}, \\ |e^{-\eta y^+} \partial_x^\alpha \mathcal{R} \ell_4^+| &\leq C e^{-\theta_1 t} (e^{-\eta|x-y|/2} + e^{-(x-y)^2/Mt}). \end{aligned}$$

Proof. By (3.62)(iii), the contribution of the reactive modes to S is

$$(3.67) \quad \chi_{\{y>0\}} \chi_{\{t \geq 1\}} \frac{e^x}{e^x + e^{-x}} r_4^+ \ell_4^{+\text{tr}} (4\pi dt)^{-1/2} e^{-(x-y+st)^2/4dt}.$$

Given $0 \leq x \leq y$, we can bound $e^{-\eta y} e^{-(x-y+st)^2/4dt}$, for $|x-y| \leq \frac{1}{2}st$, by

$$e^{-\eta y} e^{-(st/2)^2/4t} \leq e^{-\eta y} e^{-s^2 t/16} \leq e^{-\eta|y-x|} e^{-s^2 t/16},$$

and, for $|x-y| > \frac{1}{2}st$, by

$$e^{-\eta y} \leq e^{-\eta|y-x|/2} e^{-\eta y/2} \leq e^{-\eta|y-x|/2} e^{-\eta st/4},$$

and this implies (3.66)(i). To prove (3.66)(ii), we note that the contribution of the parabolic terms $t^{-\frac{1}{2}}(t+1)^{-\frac{1}{2}}e^{-(x-y)^2/Mt}$ and $\mathcal{S}_0$ to $\mathcal{R} \ell_4^+$ comes from Riemann saddle-point estimates of the sole scattered terms $\mathcal{S}_\lambda \ell_4^+$ (see the proof of Proposition 7.1 in [MaZ3] for more details). Hence (3.66)(i) implies (3.66)(ii). $\square$

Remark 3.28. The proof of Proposition 7.1 of [MaZ3] shows that Proposition 3.23 applies more generally to linear operators of the form (1.15) that satisfy (1.18) and the conditions (A1)-(A2), (H1)-(H4) of Section 1.5.

3.2.2 Convolution bounds

From the pointwise bounds of Proposition 3.23 and Remarks 3.26 and 3.27, we obtain by standard convolution bounds the following $L^p \rightarrow L^q$ estimates, exactly as described in [MaZ1, MaZ2, MaZ3, MaZ4, Z2] for the viscous shock case.

Corollary 3.29. *Under (1.18), for all $t \geq 1$, some $C > 0$, any $\eta > 0$, for any $1 \leq q \leq p$, $1 \leq p \leq +\infty$, and $f \in L^q \cap W^{1,p}$,*

$$\begin{aligned} \left| \int_{\mathbb{R}} (\mathcal{S} + \mathcal{R})(\cdot, t; y) f(y) dy \right|_{L^p} &\leq Ct^{-\frac{1}{2}(\frac{1}{q} - \frac{1}{p})} |f|_{L^q}, \\ \left| \int_{\mathbb{R}} \partial_y (\mathcal{S} + \mathcal{R})(\cdot, t; y) f(y) dy \right|_{L^p} &\leq Ct^{-\frac{1}{2}(\frac{1}{q} - \frac{1}{p}) - \frac{1}{2}} |f|_{L^q} + Ce^{-\eta t} |f|_{L^p}, \\ \left| \int_{\mathbb{R}} (\mathcal{S} + \mathcal{R})(\cdot, t; y) \ell_4^+ f(y) e^{-\theta y^+} dy \right|_{L^p} &\leq Ct^{-\frac{1}{2}(\frac{1}{q} - \frac{1}{p}) - \frac{1}{2}} |f|_{L^q} + Ce^{-\eta t} |f|_{L^p}, \\ \left| \int_{\mathbb{R}} \mathcal{H}(\cdot, t; y) f(y) dy \right|_{L^p} &\leq Ce^{-\eta t} |f|_{L^p}, \end{aligned}$$

where $y^+ := \max(y, 0)$. Likewise, for all x and all $t \geq 0$,

$$\begin{aligned} |\partial_y \mathbf{e}(\cdot, t)|_{L^p} + |\partial_t \mathbf{e}(\cdot, t)|_{L^p} &\leq Ct^{-\frac{1}{2}(1 - \frac{1}{p})}, \\ |\partial_t \partial_y \mathbf{e}(\cdot, t)|_{L^p} &\leq Ct^{-\frac{1}{2}(1 - \frac{1}{p}) - \frac{1}{2}}. \end{aligned}$$

4 Stability: Proof of Theorem 1.12

We often omit to indicate dependence on ε in the proof below. All the estimates are uniform in ε .

4.1 Linearized stability criterion

Proof of Theorem 1.12: linear case. Sufficiency of (1.18) for linearized orbital stability follows immediately by the bounds of Corollary 3.29, exactly as in the viscous shock case, setting

$$\delta(t) := \int_{\mathbb{R}} \mathbf{e}(x, t; y) U_0(y) dy$$

so that

$$U - \delta(t) \bar{U}' = \int_{\mathbb{R}} (\mathcal{H} + \mathcal{S} + \mathcal{R})(x, t; y) U_0(y) dy;$$

see [ZH, MaZ3, Z2] for further details. Necessity follows from more general spectral considerations not requiring the detailed bounds of Proposition 3.23; see the discussion of effective spectrum in [ZH, MaZ3, Z2]. The argument goes again exactly as in the viscous shock case. $\square$

4.2 Auxiliary energy estimate

Consider $\tilde{U}$ the solution of (1.2) issued from $\tilde{U}_0$, and let

$$(4.1) \quad U(x, t) := \tilde{U}(x + \delta(t), t) - \bar{U}(x).$$

Then, the following auxiliary energy estimate holds.

Lemma 4.1 (Proposition 4.15, [Z2]). *Under the hypotheses of Theorem 1.12, assume that $\tilde{U}_0 \in H^3$, and suppose that, for $0 \leq t \leq T$, the suprema of $|\dot{\delta}|$ and the H^3 norm of U each remain bounded by a sufficiently small constant. Then, for all $0 \leq t \leq T$, for some $\theta > 0$,*

$$|U(t)|_{H^3}^2 \leq C e^{-\theta t} |U(0)|_{H^3}^2 + C \int_0^t e^{-\theta(t-s)} (|U|_{L^2}^2 + |\dot{\delta}|^2)(s) ds.$$

4.3 Nonlinear stability

Proof of Theorem 1.12: nonlinear case. Let U be the perturbation variable associated with solution $\tilde{U}$ as in (4.1); by a Taylor expansion, U solves the perturbation equation

$$\partial_t U - LU = \partial_x Q_f(U, \partial_x U) + Q_r(U) + \dot{\delta}(t)(\bar{U}' + \partial_x U),$$

where the linear operator L is defined in (1.15), and

$$(4.2) \quad |Q_f| \leq C|U|(|U| + |\partial_x U|),$$

where C depends on $\|U\|_{L^\infty}$ and $\|\bar{U}\|_{W^{1,\infty}}$.

Lemma 4.2. *Under Assumptions 1.1, 1.2, if the temperature T associated with solution U satisfies $\|T\|_{L^\infty} < T_i - T_+$ (by Lemma 1.4, $0 < T_i - T_+$), then the nonlinear reactive term Q_r has the form*

$$(4.3) \quad Q_r(U) = \ell_4^+ e^{-\eta_0 x^+} q_r(U),$$

where $x^+ := \max(x, 0)$, $\eta_0 > 0$ is as in Corollary 1.6, and $q_r(U) = q_r(w, z)$ is a scalar such that

$$(4.4) \quad |q_r(U)| \leq C|U|^2,$$

where C depends on $\|U\|_{L^\infty}$ and $\|\bar{U}\|_{L^\infty}$.

Proof. We use the specific form $-k\phi(T)z\ell_4^+$ of the reactive source in (1.1), together with Taylor expansion

$$\begin{aligned} (\phi(\bar{T} + T)(\bar{z} + z) - (\phi(\bar{T})\bar{z} - (\phi'(\bar{T})T\bar{z} + \phi(\bar{T})z) \\ = \phi'(\bar{T})Tz + \phi''(\bar{T} + \beta T)T^2\bar{z}, \end{aligned}$$

for some $0 < \beta < 1$, and the fact that $\phi'(\bar{T} + T) \leq C e^{-\eta_0 x^+}$ for $|T| < T_i - T_+$, for $\eta > 0$ as in Corollary 1.6, by $\phi(T_+) = 0$ together with the property that $\phi'(T) \equiv 0$ for $T \leq T_i$ and exponential convergence of $\bar{U}(x)$ to U_+ as $x \rightarrow +\infty$. $\square$

Recalling the standard fact that $\bar{U}'$ is a stationary solution of the linearized equations (3.1), $L\bar{U}' = 0$, or

$$\int_{\mathbb{R}} \mathcal{G}(x, t; y) \bar{U}'(y) dy = e^{tL} \bar{U}'(x) = \bar{U}'(x),$$

we have by Duhamel's principle:

$$\begin{aligned} U(x, t) &= \delta(t) \bar{U}'(x) + \int_{\mathbb{R}} \mathcal{G}(x, t; y) U_0(y) dy \\ &\quad + \int_0^t \int_{\mathbb{R}} \mathcal{G}(x, t-s; y) \ell_4^+ e^{-\eta y^+} q_r(U)(y, s) dy ds \\ &\quad - \int_0^t \int_{\mathbb{R}} \partial_y \mathcal{G}(x, t-s; y) (Q_f(U, \partial_x U) + \dot{\delta} U)(y, s) dy ds. \end{aligned}$$

Defining

$$\begin{aligned} (4.5) \quad \delta(t) &= - \int_{\mathbb{R}} \mathbf{e}(y, t) U_0(y) dy \\ &\quad - \int_0^t \int_{\mathbb{R}} \mathbf{e}(y, t-s) \ell_4^+ e^{-\eta y^+} q_r(U)(y, s) dy ds \\ &\quad + \int_0^t \int_{\mathbb{R}} \partial_y \mathbf{e}(y, t-s) (Q_f(U, \partial_x U) + \dot{\delta} U)(y, s) dy ds, \end{aligned}$$

following [Z3, MaZ1, MaZ2, MaZ4], and recalling Proposition 3.23, we obtain finally the *reduced equations*:

$$\begin{aligned} (4.6) \quad U(x, t) &= \int_{\mathbb{R}} (\mathcal{H} + \mathcal{S} + \mathcal{R})(x, t; y) U_0(y) dy \\ &\quad + \int_0^t \int_{\mathbb{R}} \mathcal{H}(x, t-s; y) (\partial_y (Q_f(U, \partial_x U) + \dot{\delta} U) + \ell_4^+ e^{-\eta y^+} q_r(U)) dy ds \\ &\quad + \int_0^t \int_{\mathbb{R}} (\mathcal{S} + \mathcal{R})(x, t-s; y) \ell_4^+ e^{-\eta y^+} q_r(U) dy ds \\ &\quad - \int_0^t \int_{\mathbb{R}} \partial_y (\mathcal{S} + \mathcal{R})(x, t-s; y) (Q_f(U, \partial_x U) + \dot{\delta} U) dy ds, \end{aligned}$$

and, differentiating (4.5) with respect to t , and recalling Corollary 3.26:

$$\begin{aligned} (4.7) \quad \dot{\delta}(t) &= - \int_{\mathbb{R}} \partial_t \mathbf{e}(y, t) U_0(y) dy \\ &\quad + \int_0^t \int_{\mathbb{R}} \partial_y \partial_t \mathbf{e}(y, t-s) (Q_f(U, \partial_x U) + \dot{\delta} U)(y, s) dy ds. \end{aligned}$$

$$\begin{aligned} (4.8) \quad \delta(t) &= - \int_{\mathbb{R}} \mathbf{e}(y, t) U_0(y) dy \\ &\quad + \int_0^t \int_{\mathbb{R}} \partial_y \mathbf{e}(y, t-s) (Q_f(U, \partial_x U) + \dot{\delta} U)(y, s) dy ds, \end{aligned}$$

Define

$$\zeta(t) := \sup_{0 \leq s \leq t, 2 \leq p \leq \infty} \left(|U(\cdot, s)|_{L^p} (1+s)^{\frac{1}{2}(1-\frac{1}{p})} + |\dot{\delta}(s)|(1+s)^{\frac{1}{2}} + |\delta(s)| \right).$$

We shall establish:

Claim. *There exists $c_0 > 0$, such that, for all $t \geq 0$ for which a solution exists with ζ uniformly bounded by some fixed, sufficiently small constant, there holds*

$$\zeta(t) \leq c_0(|U_0|_{L^1 \cap H^3} + \zeta(t)^2).$$

From this result, it follows by continuous induction that, provided

$$|U_0|_{L^1 \cap H^3} < \frac{1}{4}c_0^2,$$

there holds

$$(4.9) \quad \zeta(t) \leq 2c_0|U_0|_{L^1 \cap H^3}$$

for all $t \geq 0$ such that ζ remains small. For, by standard short-time theory/local well-posedness in H^3 , and the standard principle of continuation, there exists a solution $U \in H^3$ on the open time-interval for which $|U|_{H^3}$ remains bounded, and on this interval ζ is well-defined and continuous. Now, let $[0, T)$ be the maximal interval on which $|U|_{H^3}$ remains strictly bounded by some fixed, sufficiently small constant $\delta > 0$. By Lemma 4.1, we have

$$\begin{aligned} |U(t)|_{H^3}^2 &\leq C|U(0)|_{H^3}^2 e^{-\theta t} + C \int_0^t e^{-\theta(t-\tau)} (|U|_{L^2}^2 + |\dot{\delta}|^2)(\tau) d\tau \\ &\leq C'(|U(0)|_{H^3}^2 + \zeta(t)^2)(1+t)^{-\frac{1}{2}}, \end{aligned}$$

for some $C, C', \theta > 0$, and so the solution continues so long as ζ remains small, with bound (4.9), at once yielding existence and the claimed sharp $L^p \cap H^3$ bounds, $2 \leq p \leq \infty$.

Proof of Claim. We must show that each of the quantities

$$|U|_{L^p}(1+s)^{\frac{1}{2}(1-\frac{1}{p})}, \quad |\dot{\delta}|(1+s)^{\frac{1}{2}}, \quad \text{and} \quad |\delta|$$

is separately bounded by $C(|U_0|_{L^1 \cap H^3} + \zeta(t)^2)$, for some $C > 0$, all $0 \leq s \leq t$, so long as ζ remains sufficiently small. By (4.6)–(4.7) and the triangle inequality, we have

$$\begin{aligned} |U|_{L^p} &\leq \text{I}_a + \text{I}_b + \text{I}_c + \text{I}_d, \\ |\dot{\delta}(t)| &\leq \text{II}_a + \text{II}_b, \\ |\delta(t)| &\leq \text{III}_a + \text{III}_b, \end{aligned}$$

where I_a is the L^p norm of the first integral term in the right-hand side of (4.6), I_b the second term, etc., and similarly Π_a is the modulus of the first term in the right-hand side of (4.7), etc.

We estimate each term in turn, following the approach of [MaZ1, MaZ4].

The linear term I_a satisfies bound

$$I_a \leq C|U_0|_{L^1 \cap L^p} (1+t)^{-\frac{1}{2}(1-\frac{1}{p})},$$

by Proposition 3.23 and Corollary 3.29.

Likewise, applying the bounds of Corollary 3.29, we have

$$\begin{aligned} I_b &\leq C\zeta(t)^2 \int_0^t e^{-\eta(t-s)} (1+s)^{-\frac{1}{2}} ds \\ &\leq C\zeta(t)^2 (1+t)^{-\frac{1}{2}}, \end{aligned}$$

and (taking $q = 2$ in the second estimate of Corollary 3.29)

$$\begin{aligned} I_c + I_d &\leq C \int_0^t e^{-\eta(t-s)} (|U|_{L^\infty} + |\partial_x U|_{L^\infty} + |\dot{\delta}|) |U|_{L^p}(s) ds \\ &\quad + C \int_0^t (t-s)^{-\frac{3}{4}+\frac{1}{2p}} (|U|_{L^\infty} + |\dot{\delta}|) |U|_{H^1}(s) ds \\ &\leq C\zeta(t)^2 \int_0^t e^{-\eta(t-s)} (1+s)^{-\frac{1}{2}(1-\frac{1}{p})-\frac{1}{2}} ds \\ &\quad + C\zeta(t)^2 \int_0^t (t-s)^{-\frac{3}{4}+\frac{1}{2p}} (1+s)^{-\frac{3}{4}} ds \\ &\leq C\zeta(t)^2 (1+t)^{-\frac{1}{2}(1-\frac{1}{p})}, \end{aligned}$$

$$\Pi_a \leq |\partial_t \mathbf{e}(t)|_{L^\infty} |U_0|_{L^1} \leq C|U_0|_{L^1} (1+t)^{-\frac{1}{2}}$$

and

$$\begin{aligned} \Pi_b &\leq \int_0^t |\partial_y \partial_t \mathbf{e}(t-s)|_{L^2} (|U|_{L^\infty} + |\dot{\delta}|) |U|_{H^1}(s) ds \\ &\leq C\zeta(t)^2 \int_0^t (t-s)^{-\frac{3}{4}} (1+s)^{-\frac{3}{4}} ds \\ &\leq C\zeta(t)^2 (1+t)^{-\frac{1}{2}}, \end{aligned}$$

while

$$\text{III}_a \leq |\mathbf{e}(t)|_{L_y^\infty} |U_0|_{L^1} \leq C|U_0|_{L^1}$$

and

$$\begin{aligned} \text{III}_b &\leq \int_0^t |\partial_y \mathbf{e}(t-s)|_{L^2} (|U|_{L^\infty} + |\dot{\delta}|) |U|_{H^1}(s) ds \\ &\leq C\zeta(t)^2 \int_0^t (t-s)^{-\frac{1}{4}} (1+s)^{-\frac{3}{4}} ds \\ &\leq C\zeta(t)^2. \end{aligned}$$

This completes the proof of the claim, establishing (1.19) for $p \geq 2$. The remaining bounds $1 \leq p < 2$ then follow by a bootstrap argument as described in [Z2]; we omit the details. $\square$

5 Bifurcation: Proof of Theorem 1.14

Given two Banach spaces X and Y , we denote by $\mathcal{L}(X, Y)$ the space of linear continuous applications from X to Y , and let $\mathcal{L}(X) := \mathcal{L}(X, X)$. We use (1.20) to denote weighted Sobolev spaces and norms. Let $x^+ := \max(0, x)$. Given a constant $\eta > 0$ and a weight function $\omega > 0$, define subspaces of $\mathcal{S}'(\mathbb{R})$ by

$$L_{\eta^+}^1 := \{f, e^{\eta(\cdot)^+} f \in L^1\}, \quad L_\omega^1 := \{f, \omega f \in L^1\}, \quad L_{\omega, \eta^+}^1 := \{f, \omega f \in L_{\eta^+}^1\}.$$

Definition 5.1. *Given a constant $\eta > 0$ and a weight function ω satisfying (1.21), define the Banach spaces*

$$\mathcal{B}_1, \mathcal{B}_2, X_1, X_2 \subset \mathcal{D}'(\mathbb{R}; \mathbb{C}_v^3 \times \mathbb{C}_z)$$

by

$$\begin{aligned} \mathcal{B}_1 &:= H^1, & \mathcal{B}_2 &:= H^1 \cap (\partial_x L^1 \times L_{\eta^+}^1), \\ X_1 &:= H_\omega^2, & X_2 &:= H_\omega^2 \cap (\partial_x L_\omega^1 \times L_{\omega, \eta^+}^1). \end{aligned}$$

with norms

$$\begin{aligned} \|(v, z)\|_{\mathcal{B}_1} &:= \|(v, z)\|_{H^1}, \\ \|(\partial_x v, z)\|_{\mathcal{B}_2} &:= \|(\partial_x v, z)\|_{H^1} + \|v\|_{L^1} + \|e^{\eta(\cdot)^+} z\|_{L^1}, \\ \|(v, z)\|_{X_1} &:= \|(v, z)\|_{H_\omega^2}, \\ \|(\partial_x v, z)\|_{X_2} &:= \|(\partial_x v, z)\|_{H_\omega^2} + \|\omega v\|_{L^1} + \|\omega e^{\eta(\cdot)^+} z\|_{L^1}. \end{aligned}$$

In particular, $X_2 \hookrightarrow X_1 \hookrightarrow \mathcal{B}_1$, with $\|\cdot\|_{\mathcal{B}_1} \leq \|\cdot\|_{X_1} \leq \|\cdot\|_{X_2}$, and $X_2 \hookrightarrow \mathcal{B}_2 \hookrightarrow \mathcal{B}_1$, with $\|\cdot\|_{\mathcal{B}_1} \leq \|\cdot\|_{\mathcal{B}_2} \leq \|\cdot\|_{X_2}$, and the unit ball in X_1 is closed in $\mathcal{B}_1$.

5.1 The perturbation equations

If $\tilde{U}^\varepsilon$ solves (1.2) with initial datum $\tilde{U}_{|t=0}^\varepsilon = \bar{U}^\varepsilon + U_0^\varepsilon$, then the perturbation variable $U(\varepsilon, x, t) := \tilde{U}^\varepsilon(x, t) - \bar{U}^\varepsilon(x)$ satisfies

$$(5.1) \quad \begin{cases} \partial_t U - L(\varepsilon)U = \partial_x Q_f(\varepsilon, U, \partial_x U) + Q_r(\varepsilon, U), \\ U(\varepsilon, x, 0) = U_0^\varepsilon(x). \end{cases}$$

The nonlinear term Q_f satisfies (4.2), while Q_r satisfies Lemma 4.2.

5.2 Coordinatization

Let $\varphi_{\pm}^{\varepsilon}$ be the eigenfunctions of $L(\varepsilon)$ associated with the bifurcation eigenvalues $\gamma(\varepsilon) \pm i\tau(\varepsilon)$, and let $\tilde{\varphi}_{\pm}^{\varepsilon}$ be the corresponding left eigenfunctions. We know from Section 3.1.3 that $(\gamma \pm i\tau)(\varepsilon) \in \mathbb{C} \setminus \mathcal{C}_- \cup \mathcal{C}_+$, hence $\varphi_{\pm}^{\varepsilon}$ decay exponentially at both $-\infty$ and $+\infty$, in particular, if in (1.21) θ_0 is small enough, then $\varphi_{\pm}^{\varepsilon} \in H_{\omega}^2$. Let Π be the L^2 -projection onto $\text{span}(\varphi_{\pm}^{\varepsilon})$ parallel to $\text{span}(\tilde{\varphi}_{\pm}^{\varepsilon})^{\perp}$. Decomposing

$$U = u_{11}\Re\varphi_+^{\varepsilon} + u_{12}\Im\varphi_+^{\varepsilon} + u_2, \quad U_0^{\varepsilon} = a_1\Re\varphi_+^{\varepsilon} + a_2\Im\varphi_+^{\varepsilon} + b,$$

where $u_{11}\Re\varphi_+^{\varepsilon} + u_{12}\Im\varphi_+^{\varepsilon}$ and $a_1\Re\varphi_+^{\varepsilon} + a_2\Im\varphi_+^{\varepsilon}$ belong to $\text{span}(\phi_{\pm}^{\varepsilon})$ (so that, in particular, u_{1j} and a_j are real), and coordinatizing as (u_1, u_2) , $u_1 := (u_{11}, u_{12}) \in \mathbb{R}^2$, we obtain after a brief calculation that U solves (5.1) if and only if its coordinates solve the system

$$(5.2) \quad \begin{cases} \partial_t u_1 = \begin{pmatrix} \gamma(\varepsilon) & \tau(\varepsilon) \\ -\tau(\varepsilon) & \gamma(\varepsilon) \end{pmatrix} u_1 + \Pi N(\varepsilon, u_1, u_2), \\ \partial_t u_2 = (1 - \Pi)L(\varepsilon)u_2 + (1 - \Pi)N(\varepsilon, u_1, u_2), \\ u_1|_{t=0} = a, \\ u_2|_{t=0} = b, \end{cases}$$

where

$$N(\varepsilon, u_1, u_2) := (\partial_x Q_f + Q_r)(\varepsilon, \bar{U}^{\varepsilon}, U).$$

Given $T_0 > 0$, there exist $\zeta_0 > 0$ and $C > 0$, such that, if $|a| + \|b\|_{H_{\omega}^2} < \zeta_0$, the initial value problem (5.2) possesses a unique solution $(u_1, u_2)(a, b, \varepsilon) \in C^0([0, T_0], \mathbb{R}^2 \times H_{\omega}^2)$ satisfying

$$(5.3) \quad \begin{aligned} C^{-1}|a| - C\|b\|_{H_{\omega}^2}^2 &\leq |u_1(t)| \leq C(|a| + \|b\|_{H_{\omega}^2}^2), \\ \|u_2(t)\|_{H_{\omega}^2} &\leq C(\|b\|_{H_{\omega}^2} + |a|^2), \\ \|\partial_{(a,b)}(u_1, u_2)(t)\|_{\mathcal{L}(\mathbb{R}^2 \times H^1, H^1)} &\leq C. \end{aligned}$$

(For more details on the initial value problem (5.2) and estimate (5.3), see [TZ2], Proposition 4.2.)

5.3 Poincaré return map

We express the period map $(a, b, \varepsilon) \rightarrow \hat{b} := u_2(a, b, \varepsilon, T)$ as a discrete dynamical system

$$(5.4) \quad \hat{b} = \mathbf{S}(\varepsilon, T)b + \tilde{N}(a, b, \varepsilon, T),$$

where

$$\mathbf{S}(\varepsilon, T) := e^{T(1-\Pi)L(\varepsilon)}$$

is the linearized solution operator in v and

$$\tilde{N}(a, b, \varepsilon, T) := \int_0^T \mathbf{S}(\varepsilon, T-s)(1-\Pi)N(\varepsilon, u_1, u_2)(s)ds$$

the difference between nonlinear and linear solution operators.

Evidently, periodic solutions of (5.2) with period T are fixed points of the period map (equilibria of (5.4)) or, equivalently, zeros of the displacement map

$$\Delta(a, b, \varepsilon, T) := (\mathbf{S}(\varepsilon, T) - \text{Id})b + N_2(a, b, \varepsilon, T).$$

5.4 Lyapunov-Schmidt reduction

We now carry out a nonstandard Lyapunov-Schmidt reduction following the “inverse temporal dynamics” framework of [TZ2], tailored for the situation that $\text{Id} - \mathbf{S}(\varepsilon, T)$ is not uniformly invertible, or, equivalently, the spectrum of $(1-\Pi)L(\varepsilon)$ is not bounded away from $\{j\pi/T\}_{j \in \mathbb{Z}}$. In the present situation, $(1-\Pi)L(\varepsilon)$ has both a 1-dimensional kernel (a consequence of (H4), see Section 1.5) and essential spectra accumulating at $\lambda = 0$, and no other purely imaginary spectra, so that $\text{Id} - \mathbf{S}(\varepsilon, T)$ inherits the same properties; see [TZ2] for further discussion.

Our goal, and the central point of the analysis, is to solve $\Delta(a, b, \varepsilon, T) = 0$ for b as a function of (a, ε, T) , eliminating the transverse variable and reducing to a standard planar bifurcation problem in the oscillatory variable a . A “forward” temporal dynamics technique would be to rewrite $\Delta = 0$ as a fixed point map

$$(5.5) \quad b = \mathbf{S}(\varepsilon, T)b + \tilde{N}(a, b, \varepsilon, T),$$

then to substitute for T an arbitrarily large integer multiple jT . In the strictly stable case $\Re \sigma((1-\Pi)L) \leq -\theta < 0$, $\|\mathbf{S}(\varepsilon, jT)\|_{\mathcal{L}(X_1)} < \frac{1}{2}$ for j sufficiently large. Noting that $\tilde{N}$ is quadratic in its dependency, we would have therefore contractivity of (5.5) with respect to b , yielding the desired reduction. However, in the absence of a spectral gap between $\sigma((1-\Pi)L)$ and the imaginary axis, $\|\mathbf{S}(\varepsilon, jT)\|_{\mathcal{L}(X_1)}$ does not decay, and may be always greater than unity; thus, this naive approach does not succeed.

The key idea in [TZ2] is to rewrite $\Delta = 0$ instead in “backward” form

$$(5.6) \quad b = (\text{Id} - \mathbf{S}(\varepsilon, T))^{-1} \tilde{N}(a, b, \varepsilon, T),$$

then show that $(\text{Id} - \mathbf{S}(\varepsilon, T))^{-1}$ is well-defined and bounded on $\text{Range } \tilde{N}$, thus obtaining contractivity by quadratic dependence of $\tilde{N}$. Since the right inverse $(\text{Id} - \mathbf{S}(\varepsilon, T))^{-1} \tilde{N}$ is formally given by $\sum_{j=0}^{\infty} \mathbf{S}(\varepsilon, jT) \tilde{N}$ this amounts to establishing convergence: a stability/cancellation estimate. Quite similar estimates appear in the nonlinear stability theory, where the interaction of linearized evolution $\mathbf{S}$ and nonlinear source $\tilde{N}$ are likewise crucial for decay. The formulation (5.6) can be viewed also as a “by-hand” version of the usual proof of the standard Implicit Function Theorem [TZ2].

Lemma 5.2. *Under the assumptions of Theorem 1.14, if the constant η in Definition 5.1 satisfies $\eta < \eta_0$, where η_0 was introduced in Corollary 1.6, then*

$$\tilde{N} : (a, b, \varepsilon, T) \in \mathbb{R}^2 \times X_1 \times \mathbb{R}^2 \rightarrow \tilde{N}(a, b, \varepsilon, T) \in X_2,$$

is quadratic order, and C^1 as a map from $\mathbb{R}^2 \times \mathcal{B}_1 \times \mathbb{R}^2$ to $\mathcal{B}_2$ for $\|b\|_{X_1}$ uniformly bounded, with

$$(5.7) \quad \begin{aligned} \|\tilde{N}\|_{X_2} &\leq C(|a| + \|b\|_{X_1})^2, \\ \|\partial_{(a,b)} \tilde{N}\|_{\mathcal{L}(\mathbb{R}^2 \times \mathcal{B}_1, \mathcal{B}_2)} &\leq C(|a| + \|b\|_{H^2}), \\ \|\partial_{(\varepsilon, T)} \tilde{N}\|_{\mathcal{L}(\mathbb{R}^2, \mathcal{B}_2)} &\leq C(|a| + \|b\|_{H^2})^2, \end{aligned}$$

Proof. We use the variational bounds of [TZ3] (see Propositions 5 and 6, [TZ3]) and Lemma 4.2. Note that, in (1.21), only $\omega^{-1} \in L^\infty$ and (1.21)(ii) were used at this point. $\square$

5.4.1 Pointwise cancellation estimate

We now develop the key cancellation estimates, adapting the pointwise semigroup methods of [ZH, MaZ3, Z2] to the present case.

Our starting point is the inverse Laplace transform representation (3.55). Deforming the contour using analyticity of $\mathcal{G}_\lambda$ across oscillatory eigenvalues $\lambda_\pm(\varepsilon)$ we obtain $\mathcal{G} = \tilde{\mathcal{G}} + \mathcal{O}$, where

$$\mathcal{O}(x, t; y) := e^{\lambda_+(\varepsilon)t} \varphi_+^\varepsilon(x) \tilde{\varphi}_+^\varepsilon(y)^{\text{tr}} + e^{\lambda_-(\varepsilon)t} \varphi_-^\varepsilon(x) \tilde{\varphi}_-^\varepsilon(y)^{\text{tr}}$$

is the sum of the residues of the integrand at $\lambda_\pm$ (the right- and left-eigenfunctions $\varphi_\pm^\varepsilon$ and $\tilde{\varphi}_\pm^\varepsilon$ are defined in Section 5.2). The Green function $\tilde{\mathcal{G}}$ is the kernel of the integral operator $\mathbf{S}(\varepsilon, t)$ defined in Section 5.3. Note that, under the assumptions of Theorem 1.14, the Evans function associated with $(1 - \Pi)L(\varepsilon)$ satisfies (1.18), so that, by Remark 3.28, Proposition 3.23 applies to $\tilde{\mathcal{G}}$.

For $\nu, \nu_0 > 0$, let Γ be the counterclockwise arc of circle $\partial B(0, r)$ (r as in Proposition 3.21) connecting $-\nu - i\nu_0$ and $-\nu + i\nu_0$. If ν and $\nu_0 > 0$ are sufficiently small, then Γ is entirely contained in the resolvent set of $(1 - \Pi)L(\varepsilon)$, and $\tilde{\mathcal{G}}$ can be decomposed as $\mathcal{G}_\text{I} + \mathcal{G}_\text{II}$, with

$$(5.8) \quad \begin{aligned} \mathcal{G}_\text{I}(\varepsilon, x, t; y) &:= \frac{1}{2\pi i} \int_\Gamma e^{\lambda t} \mathcal{G}_\lambda(\varepsilon, x, y) d\lambda \\ \mathcal{G}_\text{II}(\varepsilon, x, t; y) &:= \frac{1}{2\pi i} \text{P.V.} \left(\int_{-\nu-i\infty}^{-\nu-i\nu_0} + \int_{-\nu+i\nu_0}^{-\nu+i\infty} \right) e^{\lambda t} \mathcal{G}_\lambda(\varepsilon, x, y) d\lambda. \end{aligned}$$

Let $\mathbf{S}_\text{I}$ and $\mathbf{S}_\text{II}$ denote the integral operators with respective kernels $\mathcal{G}_\text{I}$ and $\mathcal{G}_\text{II}$, so that $\mathbf{S} = \mathbf{S}_\text{I} + \mathbf{S}_\text{II}$, and let $\Omega := (-\bar{\varepsilon}_0, \bar{\varepsilon}_0) \times (0, +\infty)$, for some $\bar{\varepsilon}_0 > 0$.

Remark 5.3. *The contour Γ being contained in the resolvent set of L , the elementary bound holds:*

$$|\partial_y \mathcal{G}_\lambda| \leq C e^{-\theta_\nu |x-y|}, \quad \lambda \in \Gamma,$$

for some $\theta_\nu > 0$ depending on ν . See for instance Proposition 4.4, [MaZ3].

Our treatment of the high-frequency term follows [TZ3]:

Lemma 5.4. *Under the assumptions of Theorem 1.14, the sequence of operators with kernel $\sum_{n=0}^N \mathcal{G}_{\text{II}}(\varepsilon, nT)$ is absolutely convergent in $\mathcal{L}(H^1)$, uniformly in $(\varepsilon, T) \in \Omega$.*

Proof. Starting from the description of the resolvent kernel given in Proposition 3.21, we find by the same inverse Laplace transform estimates that give terms $\mathcal{H}$ and $\mathcal{R}$ in Proposition 3.23, that the high-frequency resolvent kernel $\mathcal{G}_{\text{II}}$, defined in (5.8), may be expressed as

$$(5.9) \quad \mathcal{G}_{\text{II}} = C e^{-\theta(|x-y|+t)} + h \tau_{x+st} \delta,$$

where C and its space-time derivatives are bounded, $\theta > 0$, and $h \tau_{x+st} \delta$ is a generic hyperbolic term; in particular h has the form (3.61) and satisfies (3.64)(i). The Lemma follows. $\square$

Next we turn to the low-frequency component of $\tilde{\mathcal{G}}$. Its fluid terms are handled as in [TZ3]:

Lemma 5.5. *Under the assumptions of Theorem 1.14, the sequence of operators with kernel $\sum_{n=0}^N \mathcal{G}_{\text{I}}(\varepsilon, nT)$ converges in $\mathcal{L}(\partial_x L^1, H^1)$, uniformly with respect to $(\varepsilon, T) \in \Omega$.*

Proof. We argue as in the proof of Proposition 3 of [TZ3]. Let $f \in L^1$. By (5.8), $\sum_{n=0}^{N-1} \partial_y \mathcal{G}_{\text{I}} f$ decomposes into $\text{I} - \text{II}_N$, where

$$\begin{aligned} \text{I} &= \frac{1}{2i\pi} \int_{\mathbb{R}} \int_{\Gamma} \frac{1}{1 - e^{\lambda T}} \partial_y \mathcal{G}_\lambda f \, d\lambda \, dy, \\ \text{II}_N &= \frac{1}{2i\pi} \int_{\mathbb{R}} \int_{\Gamma} \frac{e^{NT\lambda}}{1 - e^{\lambda T}} \partial_y \mathcal{G}_\lambda f \, d\lambda \, dy. \end{aligned}$$

For small ν and $\lambda \in \Gamma$, $(1 - e^{\lambda T})^{-1} = \lambda^{-1} T^{-1} (1 + O(\lambda))$.

The boundary term I is independent of N and is seen to belong to H^1 by Remark 5.3.

By (3.56)-(3.60), $\lambda^{-1} \partial_y \mathcal{E}_\lambda$ and $\lambda^{-1} \partial_y \mathcal{S}_\lambda$ have the same form as $\mathcal{E}_\lambda$ and $\mathcal{S}_\lambda$. By Proposition 3.21, $\lambda^{-1} \partial_y \mathcal{R}_\lambda$ behaves like the sum of $\mathcal{R}_\lambda$ and a pole term of form $\lambda^{-1} e^{-\theta|x-y|}$. Hence, by the same Riemann saddle-point estimates used to bound $\mathcal{G}$ in Proposition 3.23, we find that

$$(5.10) \quad \int_{\Gamma} \frac{e^{\lambda NT}}{1 - e^{\lambda T}} \partial_y \mathcal{G}_\lambda \, d\lambda = (\mathcal{E} + \mathcal{S} + \mathcal{R})(\varepsilon, NT),$$

up to a constant (independent of N) term of the form $Ce^{-\theta|x-y|}$, where the space-time derivatives of C are uniformly bounded. This constant term satisfies the same bound as term I.

In (5.10), $\mathcal{E}$, $\mathcal{S}$, $\mathcal{R}$ denote generic excited, scattered and residual terms of form (3.58), (3.59)-(3.60) and (3.64)(ii). By dominated convergence,

$$H^1\text{-}\lim_{N \rightarrow \infty} \int_{\mathbb{R}} \mathcal{E}(\varepsilon, NT) f(y) dy$$

exists and is equal to a sum of terms of the form

$$(5.11) \quad C(\varepsilon, T)(\bar{U}^\varepsilon)' \int_{\mathbb{R}} f(y) dy.$$

Besides, by (3.59)-(3.60) and (3.64),

$$(5.12) \quad \left\| \int_{\mathbb{R}} (\mathcal{S} + \mathcal{R})(\varepsilon, NT) f(y) dy \right\|_{H^1} \leq C(NT)^{-\text{frac}14} \|f\|_{L^1}.$$

This proves convergence in H^1 of the sequence Π_N . $\square$

We examine finally the contribution to the series $\sum_n \mathbf{S}(\varepsilon, nT)$ of the new (not present in [TZ3]), reactive terms.

Lemma 5.6. *Under the assumptions of Theorem 1.14, the sequence of operators with kernels $\sum_{n=0}^N \mathcal{G}_1(\varepsilon, nT) \ell_4^+$ is absolutely convergent in $\mathcal{L}(L_{\eta^+}^1, H^1)$, uniformly with respect to $(\varepsilon, T) \in \Omega$.*

Proof. Let $f \in L^1$. By (5.8), Proposition 3.23 and Corollary 3.26, the low-frequency kernel $\mathcal{G}_1$ satisfies

$$(5.13) \quad \int_{\mathbb{R}} e^{-\eta y^+} \mathcal{G}_1(\varepsilon, t) \ell_4^+ f(y) dy = \int_{\mathbb{R}} e^{-\eta y^+} (\mathcal{S} + \mathcal{R})(\varepsilon, t) \ell_4^+ f(y) dy,$$

and, by Corollary 3.27,

$$\left\| \int_{\mathbb{R}} e^{-\eta y^+} (\mathcal{S} + \mathcal{R})(\varepsilon, t) \ell_4^+ f(y) dy \right\|_{H^1} \leq C(1 + t^{\frac{1}{4}}) e^{-\theta_1 t} \|f\|_{L^1},$$

and the upper bound defines for $t = NT$ an absolutely converging series in H^1 . $\square$

From Lemmas 5.4, 5.5 and 5.6 and the fact that $\mathbf{S}(\varepsilon, T) \in \mathcal{L}(\mathcal{B}_1)$, for all $(\varepsilon, T) \in \Omega$, we can conclude that, under the assumptions of Theorem 1.14, the operator $\text{Id} - \mathbf{S}(\varepsilon, T)$ has a right inverse

$$(\text{Id} - \mathbf{S}(\varepsilon, T))^{-1} := \sum_{n \geq 0} \mathbf{S}(\varepsilon, nT) : \quad \mathcal{B}_2 \rightarrow \mathcal{B}_1,$$

that belongs to $\mathcal{L}(\mathcal{B}_2, \mathcal{B}_1)$, locally uniformly in $(\varepsilon, T) \in \Omega$.

We will need the following regularity result for the right inverse:

Lemma 5.7. *Under the assumptions of Theorem 1.14, the operator $(\text{Id} - \mathbf{S}(\varepsilon, T))^{-1}$ is C^1 in $(\varepsilon, T) \in \Omega$, with respect to the $\mathcal{L}(\mathcal{B}_2, \mathcal{B}_1)$ norm on $\mathcal{B}_2$.*

Proof. Note that, by (3.55), $\partial_t \mathcal{G}$ has kernel $\lambda \mathcal{G}_\lambda$; in particular, the small λ (low-frequency) estimates of the proofs of Lemmas 5.5 and 5.6 imply the convergence of the sequence $\sum_{n=0}^N \partial_T \mathbf{S}_I(\varepsilon, nT)$ in $\mathcal{L}(\mathcal{B}_2, \mathcal{B}_1)$. The contribution of $\partial_T \mathcal{G}_{II}(\varepsilon, nT)$ is handled as in Lemma 5.4, by (5.9) and (3.64)(i) with $k = 1$.

Bounds for ε -derivatives are handled as in [TZ3], using either the variational equation $(L - \lambda)\partial_\varepsilon \mathcal{G}_\lambda = -(\partial_\varepsilon L)\mathcal{G}_\lambda$, or the $\partial_\varepsilon \mathcal{G}$ bounds of Proposition 3.11 from [TZ2]. $\square$

Note that the ε -derivative bound (5.7)(iii) is stated on a proper subspace of $\mathcal{B}_1$, namely X_1 . In this respect, the following Lemma, asserting boundedness of the right inverse on $X_2 \hookrightarrow \mathcal{B}_2$, in $\mathcal{L}(X_2, X_1)$ norm, is key to the reduction procedure of the following Section. (See Remark 5.12.)

Lemma 5.8. *Under the assumptions of Theorem 1.14, $(\text{Id} - \mathbf{S}(\varepsilon, T))^{-1}$ belongs to $\mathcal{L}(X_2, X_1)$, for all $(\varepsilon, T) \in \Omega$.*

Proof. The convolution bound

$$(5.14) \quad \left\| \omega^{\frac{1}{2}} \int_{\mathbb{R}} e^{-\theta|x-y|} f(y) dy \right\|_{L^2} \leq C \min(\|f\|_{L_\omega^2}, \|f\|_{L_\omega^1}),$$

where C depends on $\|\omega^{\frac{1}{2}} e^{-\theta|\cdot|}\|_{L^1 \cap L^2}$, holds by (1.21)(i) and (1.21)(iii). It implies that the contributions of $\mathcal{G}_{II}$, of I and of the constant pole term in Π_N (see the proofs of Lemmas 5.4 and 5.5) are all bounded in $\mathcal{L}(X_1)$.

The scattered and residual terms in Π contribute nothing to the limit, by (5.12).

We use again Corollary 3.27 to handle the contribution of the reactive term. In (3.66), there are two terms in the upper bound for $(\mathcal{S} + \mathcal{R})\ell_4^+$. The first term is handled by (5.14), and the second by

$$(5.15) \quad e^{-\theta t} \left\| \omega^{\frac{1}{2}} \int_{\mathbb{R}} e^{-|x-y|^2/Mt} f(y) dy \right\|_{L^2} \leq e^{-\theta t} \left\| \omega^{\frac{1}{2}} e^{-|\cdot|^2/Mt} \right\|_{L^2} \|f\|_{L_\omega^1},$$

noting that

$$(5.16) \quad \begin{aligned} e^{-\theta t} \left\| \omega^{\frac{1}{2}} e^{-|x-y|^2/Mt} \right\|_{L^2} &\leq C e^{-\theta_1 t} \left(\left\| \omega^{\frac{1}{2}} e^{-|\cdot|^2/Mt} \right\|_{L_{|x|<t}^2} + \left\| \omega^{\frac{1}{2}} e^{-|\cdot|^2/Mt} \right\|_{L_{t \leq |x|}^2} \right) \\ &\leq C e^{-\theta_1 t} \left(t^{\frac{3}{4}} \omega(t)^{\frac{1}{2}} + \left\| \omega^{\frac{1}{2}} e^{-|\cdot|^2/M} \right\|_{L^2} \right), \end{aligned}$$

and the upper bound in (5.16) defines for $t = NT$ an absolutely converging series, if θ_0 in (1.21) is small enough. We used in (5.16) the growth assumption on ω . $\square$

Remark 5.9. The above proof shows that, in (1.21), we need in particular $\theta_0 < \frac{1}{2}\eta_0$, where η_0 is the decay rate of the background profile (see Corollary 1.6). In addition, we need $\theta_0 < \min(\theta_\nu, \theta, \frac{2}{M})$, where θ is the rate of decay in the upper bounds of Proposition 3.23 and M is the diffusion rate in the upper bound of (3.66)(ii), and θ_0 to be small enough so that $\varphi_\pm^\varepsilon \in H_\omega^2$.

5.4.2 Reduction

Corollary 5.10. Under the Assumptions of Theorem 1.14, the equation

$$(5.17) \quad \Delta(a, b, \varepsilon, T) = 0 \quad (a, b, \varepsilon, T) \in \mathbb{R}^2 \times X_1 \times \mathbb{R}^2,$$

where Δ is defined in Section 5.3, is equivalent to

$$b = (\text{Id} - \mathbf{S}(\varepsilon, T))^{-1} \tilde{N}(a, b, \varepsilon, T) + \omega$$

for

$$\omega \in \text{Ker}(\text{Id} - \mathbf{S}(\varepsilon, T)) \cap X_1.$$

Proof. A simple consequence of the definition in the above Section of the right inverse $(\text{Id} - \mathbf{S}(\varepsilon, T))^{-1}$. For more details, see the proof of Lemma 2.8 in [TZ2]. $\square$

Note that, as a consequence of (H4), Section 1.5, the kernel of $\text{Id} - \mathbf{S}(\varepsilon, T)$ is of dimension one, for all ε, T , generated by $(\bar{U}^\varepsilon)'$.

Corollary 5.11. Under the Assumptions of Theorem 1.14, the map

$$\mathcal{T}(a, b, \varepsilon, T, \alpha) := (\text{Id} - \mathbf{S}(\varepsilon, T))^{-1} \tilde{N}(a, b, \varepsilon, T) + \alpha(\bar{U}^\varepsilon)',$$

is bounded from $\mathbb{R}^2 \times X_1 \times \mathbb{R}^{2+1}$ to X_1 , and C^1 from $\mathbb{R}^2 \times \mathcal{B}_1 \times \mathbb{R}^{2+1}$ to X_1 , for $|\alpha|$ bounded and $|a| + \|b\|_{X_1} + |\varepsilon|$ sufficiently small, with

$$\begin{aligned} \|\mathcal{T}\|_{X_1} &\leq C(|a| + \|b\|_{X_1}^2), \\ \|\partial_{(a,b)}\mathcal{T}\|_{\mathcal{L}(\mathbb{R}^2 \times \mathcal{B}_1, \mathcal{B}_1)} &\leq C(|a| + \|b\|_{X_1}), \\ \|\partial_T\mathcal{T}\|_{\mathcal{B}_1} &\leq C(|a|^2 + \|b\|_{X_1}^2), \\ \|\partial_\varepsilon\mathcal{T}\|_{\mathcal{B}_1} &\leq C(|a|^2 + \|b\|_{X_1}^2 + |\alpha|), \\ \|\partial_\alpha\mathcal{T}\|_{\mathcal{B}_1} &\leq C(|a|^2 + \|b\|_{X_1}^2 + 1). \end{aligned}$$

Proof. Follows from Lemma 5.2, the results of Section 5.4.1, and the above remark on the kernel of $\text{Id} - \mathbf{S}(\varepsilon, T)$. $\square$

Remark 5.12. Without Lemma 5.8 but with Lemmas 5.4 to 5.7, we could see $\mathcal{T}$ has a C^1 map from $\mathcal{B}_1$ to $\mathcal{B}_1$ (for fixed $a, \varepsilon, T, \alpha$), with quadratic bound $\|\mathcal{T}\|_{\mathcal{B}_1} \leq C\|b\|_{\mathcal{B}_1}^2$, and derivative bound $\|\partial_b\mathcal{T}\|_{\mathcal{L}(\mathcal{B}_1)} \leq C\|b\|_{\mathcal{B}_1}$. This would be sufficient to prove existence of a fixed point $b \in \mathcal{B}_1$ as in the following Proposition, but not to prove

regularity of the fixed point with respect to ε , precisely because the ε -derivative bound (5.7)(iii) requires more regularity in b , by $\partial_\varepsilon e^{tL(\varepsilon)} = e^{tL(\varepsilon)} \partial_\varepsilon L$ (see also Remark 3.12, [TZ2]). Without regularity of the fixed point of $\mathcal{T}$, we could not use the standard implicit function theorem in Section 5.4.3. These issues were discussed in detail in Section 2.3 of [TZ2].

Proposition 5.13. *Under the assumptions of Theorem 1.14, there exists a function $\beta(a, \varepsilon, T, \alpha)$, bounded from $\mathbb{R}^{4+1}$ to X_1 and C^1 from $\mathbb{R}^{4+1}$ to $\mathcal{B}_1$, such that*

$$\begin{aligned} \Delta(a, \beta(a, \varepsilon, T, \alpha), \varepsilon, T) &\equiv 0, \\ (5.18) \quad \|\beta\|_{X_1} + \|\partial_{(\varepsilon, T)} \beta\|_{\mathcal{L}(\mathbb{R}^2, \mathcal{B}_1)} &\leq C(|a|^2 + |\alpha|), \\ \|\partial_a \beta\|_{\mathcal{L}(\mathbb{R}^2, \mathcal{B}_1)} &\leq C|a|, \\ \|\partial_\alpha \beta\|_{\mathcal{L}(\mathbb{R}, \mathcal{B}_1)} &\leq C, \end{aligned}$$

for $|(a, \varepsilon, \alpha)|$ sufficiently small. Moreover, for $|(a, \varepsilon)|$, $\|b\|_{X_1}$ sufficiently small, all solutions of (5.17) lie on the 1-parameter manifold

$$\{b = \beta(a, \varepsilon, T, \alpha), \quad \alpha \in \mathbb{R}\}.$$

Proof. A consequence of the Banach fixed point theorem applied to map $\mathcal{T}$. For more details, see the proof of Proposition 2.9 in [TZ2]. $\square$

5.4.3 Bifurcation

The bifurcation analysis is straightforward now that we have reduced to a finite-dimensional problem, the only tricky point being to deal with the 1-fold multiplicity of solutions (parametrized by α).

Define to this end

$$\tilde{\beta}(a, \varepsilon, T, \hat{\alpha}) := \beta(a, \varepsilon, T, |a|\hat{\alpha}),$$

with $\hat{\alpha}$ restricted to a ball in $\mathbb{R}^1$, noting, by (5.18), that

$$\|\tilde{\beta}\|_{X_1}, \|\partial_{(a, \varepsilon, T, \hat{\alpha})} \tilde{\beta}\|_{\mathcal{L}(\mathbb{R}^4, \mathcal{B}_1)} \leq C|a|,$$

with $\tilde{\beta}$ Lipschitz in $(a, \varepsilon, T, \hat{\alpha})$ and C^1 away from $a = 0$.

Solutions (u_1, u_2) of (5.2) originating at

$$(5.19) \quad (a, b) = (a, \tilde{\beta}(a, \varepsilon, T, \hat{a})),$$

by (5.3), remain for $0 \leq t \leq T$ in a cone

$$\mathcal{C} := \{(u_1, u_2) : |u_2| \leq C_1|u_1|\},$$

$C_1 > 0$. Indeed, (5.3) implies the bound

$$\|u_2(t)\|_{X_1} \leq C|u_1(t)|, \quad \text{for } \|b\|_{X_1} \leq C_1|a|,$$

for $C_1 > 0$ small enough, for all $0 \leq t \leq T$.

Likewise, any periodic solution of (5.2) originating in $\mathcal{C}$, since it necessarily satisfies $\Delta = 0$, must originate from data (a, b) of the form (5.19).

Defining $b \equiv \tilde{\beta}(a, \varepsilon, T, \hat{\alpha})$, and recalling invariance of $\mathcal{C}$ under flow (5.2), we may view $v(t)$ as a multiple

$$(5.20) \quad u_2(x, t) = c(a, \varepsilon, T, \hat{\alpha}, x, t)u_1(t)$$

of $u_1(t)$, where c is bounded, Lipschitz in all arguments, and C^1 away from $a = 0$. Substituting into (5.2)(i), we obtain a planar ODE

$$\partial_t u_1 = \begin{pmatrix} \gamma(\varepsilon) & \tau(\varepsilon) \\ -\tau(\varepsilon) & \gamma(\varepsilon) \end{pmatrix} u_1 + M(u_1, \varepsilon, T, t, \hat{\alpha}, a)$$

in approximate Hopf normal form, with nonlinearity $M := \Pi N$ now nonautonomous and depending on the additional parameters $(T, \hat{\alpha}, a)$, but, by (4.2) and (4.4), still satisfying the key bounds

$$(5.21) \quad |M|, |\partial_{\varepsilon, T, \hat{\alpha}} M| \leq C|u_1|^2; \quad |\partial_{a, w} M| \leq C|u_1|$$

along with planar bifurcation criterion (1.22). From (5.21), we find that M is C^1 in all arguments, also at $a = 0$. By standard arguments (see, e.g., [HK, TZ1]), we thus obtain a classical Hopf bifurcation in the variable u_1 with regularity C^1 , yielding existence and uniqueness up to time-translates of a 1-parameter family of solutions originating in $\mathcal{C}$, indexed by r and δ with $r := a_1$ and (without loss of generality) $a_2 \equiv 0$. Bound (1.23) is a consequence of (5.3)(i) and (5.20).

Finally, in order to establish uniqueness up to spatial translates, we observe, first, that, by dimensional considerations, the one-parameter family constructed must agree with the one-parameter family of spatial translates, and second, we argue as in [TZ2] that any periodic solution has a spatial translate originating in $\mathcal{C}$, yielding uniqueness up to translation among all solutions and not only those originating in $\mathcal{C}$; see Proposition 2.20 and Corollary 2.21 of [TZ2] for further details.

6 Nonlinear instability: Proof of Theorem 1.15

We describe a nonlinear instability result in general setting. Consider

$$(6.1) \quad \partial_t U = LU + \partial_x N(U) + R(U),$$

well-posed in H^s , where

$$L = \partial_x(B\partial_x U) + \partial_x(AU) + GU,$$

and $|N(U)|, |R(U)| \leq C|U|^2$ for $|U| \leq C$. Suppose that L has a conjugate pair of simple unstable eigenvalues $\lambda_{\pm} = \gamma \pm i\tau$, $\gamma > 0$, and the rest of the spectrum is neutrally stable, without loss of generality $e^{(1-\Pi)Lt} \leq Ct$, where Π is the projection onto the eigenspace associated with $\lambda_{\pm}$.

Coordinatizing similarly as in Section 5 by

$$U(x, t) = u_{11}\varphi_1(x) + u_{12}\varphi_2(x) + u_2(x, t),$$

where $\varphi_j = O(e^{-\theta|x|})$ are eigenfunctions of L , denote $r(t) := |u_1|(t)$. Then, so long as $|U|_{H^s} \leq C\epsilon$, we have existence (by variation of constants, standard continuation) of solutions of (6.1) in H^s , with estimates

$$(6.2) \quad \begin{aligned} r' &= \gamma r + O(\epsilon)|U|, \\ u_2' &= (1 - \Pi)Lu_2 + O(\epsilon)|U| \end{aligned}$$

in L^2 .

We shall argue by contradiction. That is, using (6.2), we shall show, for $C > 0$ fixed, $\epsilon > 0$ sufficiently small, and $|u_2(0)|_{H^1} \leq Cr(0)$, that eventually $r(t) \geq \epsilon$, no matter how small is $r(0)$, or equivalently $|U|_{H^s}(0)$. This, of course entails nonlinear instability.

Define $\alpha(x, t) := u_2(x, t)/r(t)$. Then,

$$\alpha' = \frac{ru_2' - u_2r'}{r^2} = \frac{u_2'}{r} - \frac{u_2}{r} \frac{r'}{r},$$

yielding after some rearrangement the equation

$$(6.3) \quad \alpha' = ((1 - \Pi)L - \gamma)\alpha + O(\epsilon(e^{-\theta|x|} + |\alpha| + |\alpha|^2)).$$

From (6.3) and standard variation of constants/contraction mapping argument, we find that $|\alpha(t)|_{H^1}$ remains less than or equal to $C|\alpha(t_0)|_{H^1}$ for $t - t_0$ small.

By variation of constants and the semigroup bound $|e^{((1-\Pi)L-\gamma)t}|_{H^1 \rightarrow H^1} \leq Ce^{-\gamma t}$ (note: γ is scalar so commutes with $(1 - \Pi)L$), we obtain

$$\delta(t) \leq C(|\alpha(0)|_{H^1} + \epsilon(1 + \delta(t)^2)),$$

for $\delta(t) := \sup_{0 \leq \tau \leq t} |\alpha(\tau)|_{H^1}$. So long as δ remains less than or equal to unity and $C\epsilon \leq \frac{1}{2}$, this yields

$$\delta(t) \leq 2C(|\alpha(0)|_{H^1} + \epsilon),$$

and thus

$$\delta(t) \leq 2C|\alpha(0)|_{H^1} + \frac{1}{2}.$$

Substituting into the radial equation, we obtain

$$r' \geq (\gamma - (1 + \delta)\epsilon)r,$$

yielding exponential growth for ϵ sufficiently small. In particular, $r \geq C\epsilon$ for some time, and thus $|U|_{H^1} \geq \epsilon$, a contradiction. We may conclude, therefore, that $|U|_{L^2}$ eventually grows larger than any ϵ , no matter how small the initial size $r(0)$, and thus we may conclude *instability* of the trivial solution $U \equiv 0$. Taking now (6.1) to be the perturbation equations about a strong detonation profile, we obtain the result of nonlinear instability of the background profile $\bar{U}$.

Remark 6.1. *In the easier case of a single, real eigenvalue, the scalar, w equation, would play the role of the radial equation here. This case is subsumed in our analysis as well.*

References

- [AT] G. Abouseif and T.Y. Toong, *Theory of unstable one-dimensional detonations*, Combust. Flame 45 (1982) 67–94.
- [AGJ] J. Alexander, R. Gardner and C.K.R.T. Jones, *A topological invariant arising in the analysis of traveling waves*, J. Reine Angew. Math. 410 (1990) 167–212.
- [AIT] R. L. Alpert and T.Y. Toong, *Periodicity in exothermic hypersonic flows about blunt projectiles*, Acta Astron. 17 (1972) 538–560.
- [BHRZ] B. Barker, J. Humpherys, K. Rudd, and K. Zumbrun, *Stability of viscous shocks in isentropic gas dynamics*, Comm. Math. Phys. 281 (2008), no. 1, 231–249.
- [Ba] G.K. Batchelor, *An introduction to fluid dynamics*, Second paperback edition. Cambridge Mathematical Library. Cambridge University Press, Cambridge (1999) xviii+615 pp. ISBN: 0-521-66396-2.
- [BeSZ] M. Beck, B. Sandstede, and K. Zumbrun, *Nonlinear stability of time-periodic shock waves*, preprint (2008).
- [BMR] A. Bourlioux, A. Majda, and V. Roytburd, *Theoretical and numerical structure for unstable one-dimensional detonations*. SIAM J. Appl. Math. 51 (1991) 303–343.
- [BDG] T.J. Bridges, G. Derks, and G. Gottwald, *Stability and instability of solitary waves of the fifth-order KdV equation: a numerical framework*. Phys. D 172 (2002), no. 1-4, 190–216.
- [Br1] L. Brin, *Numerical testing of the stability of viscous shock waves*. Doctoral thesis, Indiana University (1998).
- [Br2] L. Q. Brin, *Numerical testing of the stability of viscous shock waves*. Math. Comp. 70 (2001) 235, 1071–1088.
- [BrZ] L. Brin and K. Zumbrun, *Analytically varying eigenvectors and the stability of viscous shock waves*. Seventh Workshop on Partial Differential Equations, Part I (Rio de Janeiro, 2001). Mat. Contemp. 22 (2002), 19–32.
- [B] J.D. Buckmaster, *An introduction to combustion theory*, 3–46, in *The mathematics of combustion*, Frontiers in App. Math. (1985) SIAM, Philadelphia ISBN: 0-89871-053-7.

- [C] J. Carr, *Applications of centre manifold theory*. Applied Mathematical Sciences, 35. Springer-Verlag, New York-Berlin, 1981. vi+142 pp. ISBN: 0-387-90577-4.
- [Ch] G. Q. Chen, *Global solutions to the compressible Navier-Stokes equations for a reacting mixture*. SIAM J. Math. Anal. 23 (1992), no. 3, 609–634.
- [CF] R. Courant and K.O. Friedrichs, *Supersonic flow and shock waves*, Springer-Verlag, New York (1976) xvi+464 pp.
- [EE] D.E. Edmunds and W.D. Evans, *Spectral theory and differential operators*. Oxford University Press (1987).
- [Er] J. J. Erpenbeck, *Nonlinear theory of unstable one-dimensional detonations*, Phys. Fluids 10 (1967) No. 2, 274–289.
- [F1] W. Fickett, *Stability of the square wave detonation in a model system*. Physica 16D (1985) 358–370.
- [F2] W. Fickett, *Detonation in miniature*, 133–182, in *The mathematics of combustion*, Frontiers in App. Math. (1985) SIAM, Philadelphia ISBN: 0-89871-053-7.
- [FW] Fickett and Wood, *Flow calculations for pulsating one-dimensional detonations*. Phys. Fluids 9 (1966) 903–916.
- [G] R. Gardner, *On the detonation of a combustible gas*, Trans. Amer. Math. Soc. 277 (1983), no. 2, 431–468.
- [GK] I. Gohberg and M.G. Krein, *Introduction to the theory of linear nonselfadjoint operators*. American Mathematical Society. Translations of mathematical monographs volume 18 (1969).
- [GZ] R. Gardner and K. Zumbrun, *The Gap Lemma and geometric criteria for instability of viscous shock profiles*, Comm. Pure Appl. Math. 51 (1998), no. 7, 797–855.
- [GS1] I. Gasser and P. Szmolyan, *A geometric singular perturbation analysis of detonation and deflagration waves*, SIAM J. Math. Anal. 24 (1993) 968–986.
- [GS2] I. Gasser and P. Szmolyan, *Detonation and deflagration waves with multistep reaction schemes*, SIAM J. Appl. Math. 55 (1995) 175–191.
- [HK] J. Hale and H. Koçak, *Dynamics and bifurcations*, Texts in Applied Mathematics, 3. Springer-Verlag, New York, 1991. xiv+568 pp. ISBN: 0-387-97141-6.
- [He] D. Henry, *Geometric theory of semilinear parabolic equations*. Lecture Notes in Mathematics volume 840, Springer-Verlag, Berlin (1981), iv + 348 pp.
- [HZ] P. Howard and K. Zumbrun, *Stability of undercompressive viscous shock waves*, in press, J. Differential Equations 225 (2006), no. 1, 308–360.
- [HuZ] J. Humpherys and K. Zumbrun, *An efficient shooting algorithm for Evans function calculations in large systems*, Phys. D 220 (2006), no. 2, 116–126.
- [JLW] H.K. Jenssen, G. Lyng, and M. Williams, *Equivalence of low-frequency stability conditions for multidimensional detonations in three models of combustion*, Indiana Univ. Math. J. 54 (2005), no. 1, 1–64.

- [Kat] T. Kato, *Perturbation theory for linear operators*. Springer–Verlag, Berlin Heidelberg (1985).
- [KS] A.R. Kasimov and D.S. Stewart, *Spinning instability of gaseous detonations*. J. Fluid Mech. 466 (2002), 179–203.
- [LS] *Calculation of linear detonation instability: one-dimensional instability of plane detonation*, J. Fluid Mech. 216 (1990) 102–132.
- [LyZ1] G. Lyng and K. Zumbrun, *A stability index for detonation waves in Majda’s model for reacting flow*. Phys. D 194 (2004), no. 1-2, 1–29.
- [LyZ2] G. Lyng and K. Zumbrun, *One-dimensional stability of viscous strong detonation waves*. Arch. Ration. Mech. Anal. 173 (2004), no. 2, 213–277.
- [LRTZ] G. Lyng, M. Raoofi, B. Texier, and K. Zumbrun, *Pointwise Green Function Bounds and stability of combustion waves*, to appear, J. Diff. Eq. (2007).
- [MM] J. E. Marsden, M. McCracken, *The Hopf bifurcation and its applications*, Applied Mathematical Sciences 19, Springer.
- [MaZ1] C. Mascia and K. Zumbrun, *Pointwise Green’s function bounds and stability of relaxation shocks*. Indiana Univ. Math. J. 51 (2002), no. 4, 773–904.
- [MaZ2] C. Mascia and K. Zumbrun, *Stability of small-amplitude shock profiles of symmetric hyperbolic-parabolic systems*, Comm. Pure Appl. Math. 57 (2004), no. 7, 841–876.
- [MaZ3] C. Mascia and K. Zumbrun, *Pointwise Green function bounds for shock profiles of systems with real viscosity*, Arch. Ration. Mech. Anal. 169 (2003), no. 3, 177–263.
- [MaZ4] C. Mascia and K. Zumbrun, *Stability of large-amplitude viscous shock profiles of hyperbolic-parabolic systems*, Arch. Ration. Mech. Anal. 172 (2004), no. 1, 93–131.
- [MaZ5] C. Mascia and K. Zumbrun, *Stability of large-amplitude shock profiles of general relaxation systems*, SIAM J. Math. Anal. 37 (2005), no. 3, 889–913.
- [MeZ] G. Métivier and K. Zumbrun, *Large viscous boundary layers for noncharacteristic nonlinear hyperbolic problems*, Mem. Amer. Math. Soc. 175 (2005), no. 826, vi+107 pp.
- [Pa] A. Pazy, *Semigroups of linear operators and applications to partial differential equations*. Applied Mathematical Sciences, 44. Springer-Verlag, New York, 1983. viii+279 pp.
- [MT] U.B. McVey and T.Y. Toong, *Mechanism of instabilities in exothermic blunt-body flows*, Combust. Sci. Tech. 3 (1971) 63–76.
- [RZ] R. Raoofi and K. Zumbrun, *Stability of undercompressive viscous shock profiles of hyperbolic-parabolic systems*, Preprint (2007).
- [SS] B. Sandstede and A. Scheel, *Hopf bifurcation from viscous shock waves*, SIAM J. Math. Anal. 39 (2008), no. 6, 2033–2052.
- [ShK] S. Shizuta and Y. Kawashima, *On the normal form of the symmetric hyperbolic-parabolic systems associated with the conservation laws*. Tohoku Math. J. (2) 40 (1988), no. 3, 449–464.

- [TZ1] B. Texier and K. Zumbrun, *Relative Poincaré–Hopf bifurcation and galloping instability of traveling waves*, Methods Appl. Anal. 12 (2005), no. 4, 349–380.
- [TZ2] B. Texier and K. Zumbrun, *Galloping instability of viscous shock waves*. Physica D 237 (2008), no. 10–12, 1553–1601.
- [TZ3] B. Texier and K. Zumbrun, *Hopf bifurcation of viscous shock waves in compressible gas-dynamics and MHD*. Archive for Rational Mechanics and Analysis 190 (2008), no. 1, 107–140.
- [VT] A. Vanderbauwhede and G. Iooss, *Center manifold theory in infinite dimensions*, Dynamics reported: expositions in dynamical systems, 125–163, Dynam. Report. Expositions Dynam. Systems (N.S.), 1, Springer, Berlin, 1992.
- [Z1] K. Zumbrun, *Multidimensional stability of planar viscous shock waves*. Advances in the theory of shock waves, 307–516, Progr. Nonlinear Differential Equations Appl., 47, Birkhäuser Boston, Boston, MA, 2001.
- [Z2] K. Zumbrun, *Stability of large-amplitude shock waves of compressible Navier–Stokes equations*. Handbook of mathematical fluid dynamics. Vol. III, 311–533, North-Holland, Amsterdam, 2004.
- [Z3] K. Zumbrun, *Planar stability criteria for viscous shock waves of systems with real viscosity*, Hyperbolic systems of balance laws, 229–326, Lecture Notes in Math., 1911, Springer, Berlin, 2007.
- [ZH] K. Zumbrun and P. Howard, *Pointwise semigroup methods and stability of viscous shock waves*. Indiana Mathematics Journal V47 (1998), 741–871; Errata, Indiana Univ. Math. J. 51 (2002), no. 4, 1017–1021.
- [ZS] K. Zumbrun and D. Serre, *Viscous and inviscid stability of multidimensional planar shock fronts*, Indiana Univ. Math. J. 48 (1999) 937–992.